



# Quick Reference Guide

## HEDIS® MY 2026

FOR MORE INFORMATION, VISIT **WWW.NCQA.ORG**



Meridian, Wellcare, and Ambetter from Meridian are affiliated products serving Medicaid, Medicare, and Health Insurance Marketplace members respectively. The information presented here is representative of our network of products. If you have any questions, please contact Provider Relations.

HEDIS® is a registered trademark of the National Committee for Quality Assurance (“NCQA”). The HEDIS measures and specifications were developed by and are owned by NCQA. NCQA holds a copyright in these materials and may rescind or alter these materials at any time. Users of the HEDIS measures and specifications shall not have the right to alter, enhance or otherwise modify the HEDIS measures and specifications, and shall not disassemble, recompile or reverse engineer the HEDIS measures and specifications. Anyone desiring to use or reproduce the materials, subject to licensed user restrictions, without modification for an internal non-commercial purpose may do so without obtaining any approval from NCQA. Use of the Rules for Allowable Adjustments of HEDIS to make permitted adjustments of the materials does not constitute a modification. All other uses, including commercial use, or any external reproduction, distribution, or publication must be approved by NCQA and are subject to a license at the discretion of NCQA.

This guide has been updated with information from the release of the HEDIS® MY 2026 Volume 2 Technical Specifications by NCQA and is subject to change.

# HEDIS® MY 2026 Quick Reference Guide

**Updated to reflect NCQA HEDIS® MY 2026 Technical Specifications**

Meridian, WellCare, and Ambetter from Meridian strive to provide quality healthcare to our membership as measured through HEDIS® quality metrics. We created the HEDIS® MY 2026 Quick Reference Guide to help you increase your practice's HEDIS® rates and to use to address care opportunities for your patients. Please always follow the State and/or CMS billing guidance and ensure the HEDIS® codes are covered prior to submission.

---

## What is HEDIS®?

 HEDIS® (Healthcare Effectiveness Data and Information Set) is a set of standardized performance measures developed by the National Committee for Quality Assurance (NCQA) to objectively measure, report, and compare quality across health plans. NCQA develops HEDIS® measures through a committee represented by purchasers, consumers, health plans, health care providers, and policy makers.

## What are the scores used for?

 As state and federal governments move toward a quality-driven healthcare industry, HEDIS® rates are becoming more important for both health plans and individual providers. State purchasers of healthcare use aggregated HEDIS® rates to evaluate health insurance companies' efforts to improve preventive health outreach for members.

Physician-specific scores are also used to measure your practice's preventive care efforts. Your practice's HEDIS® score determines your rates for physician incentive programs that pay you an increased premium — for example Pay for Performance or Quality Bonus Funds.

To learn more about the Provider Incentive Programs offered by Meridian, Wellcare, and Ambetter from Meridian, please refer to the *2026 Provider Incentive Guidebook*.



# How can I submit data to improve my HEDIS® scores?

## ✓ **Administrative claims**

- Submit claim/encounter data for each service rendered with appropriate billing codes
- Submit applicable codes when members should be excluded from HEDIS measures
- Accurate and timely submission of claims data is the most appropriate way to ensure all services are captured and accounted for

## ✓ **Supplemental Data**

### • **EDI Feeds**

- Partner with our Supplemental Data team to get an Electronic Data Interchange (EDI) feed set up. This electronic method of data submission moves information directly from one computer application to another in a standard electronic format
- For information regarding set up and applicable HEDIS measures, please reach out to [katherine.g.hodges@mimeridian.com](mailto:katherine.g.hodges@mimeridian.com)

### • **EMR Access**

- Partner with our HEDIS Abstraction team to set up remote access to your EMR system for HEDIS gap closure. This access is utilized year-round to abstract and enter data for health plan prioritized HEDIS measures
- For information regarding set up and applicable HEDIS measures, please reach out to [mihedis@mhplan.com](mailto:mihedis@mhplan.com).

### • **Medical Record Submission**

- Fax medical record documentation to 833-667-1532
- Email medical record documentation to [mihedis@mhplan.com](mailto:mihedis@mhplan.com)





## Questions?



**mimeridian.com** or **mmp.mimeridian.com**



**833-667-1532**



**mihedis@mhplan.com**

Providers and other healthcare staff should document to the highest specificity to aid with the most correct coding choice.

**Ancillary staff:** Please check the tabular list for the most specific ICD-10 code choice.



For more information, visit [www.ncqa.org](http://www.ncqa.org)

## Key Changes to HEDIS Measures

- **Adult Immunization Status (AIS-E)**

Covid-19 vaccine requirement added for members 65 and older

- **Asthma Medication Ratio (AMR)**

Measure retired

- **Lead Screening in Children (LSC-E)**

Measure to be reported via Electronic Clinical Data Systems

- **Statin Therapy for Patients with Cardiovascular Disease (SPC-E)**

Measure to be reported via Electronic Clinical Data Systems

- **Statin Therapy for Patients with Diabetes (SPD-E)**

Measure to be reported via Electronic Clinical Data Systems

# Contents

## Adult Health

(AAP) Adults' Access to Preventive/Ambulatory Health Services .....	9
(ACP) Advance Care Planning .....	10
(AIS-E) Adult Immunization Status .....	12
(BPD) Blood Pressure Control for Patients With Diabetes .....	16
(CBP) Controlling High Blood Pressure .....	18
(COA) Care for Older Adults .....	20
(COL-E) Colorectal Cancer Screening .....	22
(EED) Eye Exams for Patients With Diabetes .....	25
(FMC) Follow-Up After Emergency Department Visit for People With Multiple High-Risk Chronic Conditions .....	27
(GSD) Glycemic Status Assessment for Patients With Diabetes .....	31
(KED) Kidney Health Evaluation for Patients With Diabetes .....	33
(PCE) Pharmacotherapy Management for COPD Exacerbation .....	35
(PCR) Plan All-Cause Readmissions .....	37
(SSD) Diabetes Screening for People with Schizophrenia or Bipolar Disorder Who are Using Antipsychotic Medication .....	39
(TRC) Transitions of Care .....	41

## Women's Health

(BCS-E) Breast Cancer Screening .....	44
(CCS-E) Cervical Cancer Screening .....	46
(CHL) Chlamydia Screening .....	48
(OMW) Osteoporosis Management in Women Who Had a Fracture .....	50
(OSW) Osteoporosis Screening in Older Women .....	52
(PPC) Prenatal and Postpartum Care .....	54
(PRS-E) Prenatal Immunization Status .....	58
(PND-E) Prenatal Depression Screening and Follow-Up .....	60
(PDS-E) Postpartum Depression Screening and Follow-Up .....	65

# Pediatric Health

(ADD-E) Follow-Up Care for Children Prescribed ADHD Medication .....	71
(APM-E) Metabolic Monitoring for Children and Adolescents on Antipsychotics .....	73
(CIS-E) Childhood Immunization Status.....	75
(IMA-E) Immunizations for Adolescents.....	80
(LSC-E) Lead Screening in Children .....	83
(OED) Oral Evaluation, Dental Services .....	85
(TFC) Topical Fluoride for Children.....	86
(WCV) Child and Adolescent Well-Care Visits.....	88
(W30) Well-Child Visits in the First 30 Months of Life .....	89
(WCC) Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents .....	91

# General Health

(CWP) Appropriate Testing for Pharyngitis .....	94
(FUA) Follow-Up After Emergency Department Visit for Substance Use .....	95
(FUI) Follow-Up After High-Intensity Care for Substance Use Disorder .....	100
(FUH) Follow-Up After Hospitalization for Mental Illness.....	104
(FUM) Follow-Up After Emergency Department Visit for Mental Illness.....	107
(IET) Initiation and Engagement of Substance Use Disorder Treatment .....	110
(MI2.6) Timely Transmission of Care Transition Record to Health Care Professional .....	113
(MI5.6) Medication Review .....	115
(MI7.3) Annual Dental Visit/Adults: Any Dental Visit .....	116
(URI) Appropriate Treatment for Upper Respiratory Infection.....	120
(SNS-E) Social Need Screening and Intervention .....	121

# Medication Adherence

(PDC-DIA) Medication Adherence for Diabetes Medications .....	132
(PDC-RAS) Medication Adherence for Hypertension-RAS antagonists .....	133
(PDC-STA) Medication Adherence for Cholesterol (Statins) .....	135
(SPC-E) Statin Therapy for Patients with Cardiovascular Disease .....	136
(SPD-E) Statin Therapy for Patients with Diabetes .....	138

# Adult Health

---



## (AAP) Adults' Access to Preventive/Ambulatory Health Services

### Product Lines:

- ✓ Meridian (Medicaid)
- ✓ MI Coordinated Health (Dual Eligible Plan)
- ✓ WellCare (Medicare)

### Description

Measure evaluates the percentage of members 20 years and older who had an ambulatory or preventive care visit during the measurement year.

Description	Codes*
Ambulatory Visits	<b>CPT:</b> 92002, 92004, 92012, 92014, 98966-98968, 98970-98972, 98980, 98981, 99202-99205, 99211-99215, 99242-99245, 99304-99310, 99315, 99316, 99341-99345, 99347-99350, 99381-99387, 99391-99397, 99401-99404, 99411, 99412, 99421-99423, 99429, 99441-99443, 99457, 99458, 99483 <b>HCPCS:</b> G0071, G0402, G0438, G0439, G0463, G2010, G2012, G2250, G2251, G2252, S0620, S0621, T1015
Reason for Ambulatory Visit	Z00.00, Z00.01, Z00.121, Z00.129, Z00.3, Z00.5, Z00.8, Z02.0, Z02.1, Z02.2, Z02.3, Z02.4, Z02.5, Z02.6, Z02.71, Z02.79, Z02.81, Z02.82, Z02.83, Z02.84, Z02.89, Z02.9, Z76.1, Z76.2
Hospice Care	<b>HCPCS:</b> G9473-G9479, Q5003-Q5008, Q5010, S9126, T2042-T2046, G0182 <b>CPT:</b> 99377, 99378

\*Codes subject to change

(continued)



## (AAP) Adults' Access to Preventive/Ambulatory Health Services (continued)

### Action

- ✓ Complete an appointment with all assigned patients annually

### Tips and Best Practices

- ✓ Address all preventive health screenings and tests at appropriate intervals
- ✓ Schedule next appointment at close of visit and offer flexible appointment availability, including evening, weekends, telehealth, and family appointments
- ✓ Provide appointment reminders

### Required Exclusions

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above or medical record documentation:

- ✓ Members who use hospice services any time during the measurement year
- ✓ Members who die any time during the measurement year



## (ACP) Advance Care Planning

### Product Lines:

- ✓ MI Coordinated Health (Dual Eligible Plan)
- ✓ WellCare (Medicare)

### Description

Measure evaluates percentage of adults 66-80 years of age with advanced illness, an indication of frailty or who are receiving palliative care, and adults 81 years of age and older who had advance care planning during the measurement year.

(continued)



## (ACP) Advance Care Planning (continued)

Description	Codes*
Advance Care Planning	<b>CPT:</b> 99483, 99497 <b>CPT-CAT-II:</b> 1123F, 1124F, 1157F, 1158F <b>HCPCS:</b> S0257 <b>ICD10CM:</b> Z66
Hospice Care	<b>HCPCS:</b> G9473-G9479, Q5003-Q5008, Q5010, S9126, T2042-T2046, G0182 <b>CPT:</b> 99377, 99378

\*Codes subject to change

### Action

- ✓ Talk with patients about their decisions for resuscitation, life sustaining treatment, and end-of life-care
- ✓ Even if advance directive is already on file, it must be reviewed, addressed and updated every year

### Tips and Best Practices

- ✓ Regularly encourage all patients to think about what goals they would have for care if they faced a life-threatening illness or injury; identify someone they would want to have decisions on their behalf if they did not have decision-making capacity; make their views known to their designated surrogate and to other family members
- ✓ Incorporate notes from the advance care planning discussion into the medical record
- ✓ Periodically review with the patient his or her goals, preferences, and chosen decision maker, which often change over time or with changes in health status
- ✓ Telephone visits, e-visits, or virtual check-ins are acceptable

(continued)



## (ACP) Advance Care Planning (continued)

### Required Exclusions

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above or medical record documentation:

- ✓ Members who use hospice services any time during the measurement year
- ✓ Members who die any time during the measurement year

---



## (AIS-E) Adult Immunization Status

### Product Lines:

- ✓ Meridian (Medicaid)
- ✓ MI Coordinated Health (Dual Eligible Plan)
- ✓ Ambetter from Meridian (Marketplace)
- ✓ WellCare (Medicare)

### Description

Measure evaluates the percentage of members 19 years of age and older who are up to date on recommended routine vaccines for influenza, tetanus and diphtheria (Td) or tetanus, diphtheria and acellular pertussis (Tdap), zoster, pneumococcal, hepatitis B and coronavirus disease 2019 (COVID-19).

(continued)



## (AIS-E) Adult Immunization Status (continued)

Description	Codes*
Adult Influenza Immunization	<b>CVX:</b> 88, 135, 140, 141, 144, 150, 153, 155, 158, 166, 168, 171, 185, 186, 197, 205, 111, 149 <b>CPT:</b> 90653, 90656, 90658, 90661, 90662, 90673, 90674, 90682, 90686, 90688, 90689, 90694, 90756, 90660, 90672
Anaphylaxis Due To Influenza Vaccine	<b>SNOMED:</b> 471361000124100
Td Immunization	<b>CVX:</b> 09, 113, 115, 138, 139 <b>CPT:</b> 90714
Tdap Vaccine	<b>CPT:</b> 90715
Anaphylaxis Due to Diphtheria, Tetanus or Pertussis Vaccine	<b>SNOMED:</b> 428281000124107, 428291000124105
Encephalitis Due to Diphtheria, Tetanus or Pertussis Vaccine	<b>SNOMED:</b> 192710009, 192711008, 192712001
Herpes Zoster Vaccine	<b>CVX:</b> 187 <b>CPT:</b> 90750
Anaphylaxis Due to Herpes Zoster Vaccine	<b>SNOMED:</b> 471371000124107, 471381000124105
Adult Pneumococcal Immunization	<b>CVX:</b> 33, 109, 133, 152, 215, 216, 327 <b>CPT:</b> 90670, 90671, 90677, 90684, 90732 <b>HCPCS:</b> G0009
Anaphylaxis Due to Pneumococcal Immunization	<b>SNOMED:</b> 471141000124102
Childhood Hepatitis B Immunization	<b>CVX:</b> 08, 44, 45, 51, 110, 146, 198 <b>CPT:</b> 90697, 90723, 90740, 90744, 90747, 90748 <b>HCPCS:</b> G0010
Adult Hepatitis B Vaccine (2 dose)	<b>CVX:</b> 189 <b>CPT:</b> 90739, 90743

(continued)



## (AIS-E) Adult Immunization Status (continued)

Adult Hepatitis B Vaccine (3 dose)	<b>CPT:</b> 90740, 90744, 90746, 90747, 90759
History of Hepatitis B Illness:	<b>ICD10CM:</b> B16.0, B16.1, B16.2, B16.9, B17.0, B18.0, B18.1, B19.10, B19.11
COVID-19 Vaccine	<b>CVX:</b> 309, 312, 313 <b>CPT:</b> 91304, 91320, 91322
Anaphylaxis due to COVID-19 Vaccine	<b>SNOMED:</b> 914587451000119107
Hospice Care	<b>HCPCS:</b> G9473-G9479, Q5003-Q5008, Q5010, S9126, T2042-T2046, G0182 <b>CPT:</b> 99377, 99378

\*Codes subject to change

### Action

Perform the following immunizations in the listed timeframe:

✓ Hepatitis B

Any of the following are acceptable:

- 3 doses of childhood Hep B vaccine on or before 19th birthday
- 2 doses of recommended two-dose adult Hep B vaccine after 19th birthday
- 3 doses of any other recommended adult Hep B vaccine after 19th birthday
- Positive result on Hep B testing

✓ Influenza

- Annually

✓ Td/Tdap

- Every ten years

✓ Pneumococcal

- One dose, on or after patient's 19th birthday

✓ Zoster

- Two doses, both completed on or after October 20, 2017

✓ COVID-19

- Annually, on or after 65th birthday

(continued)



## (AIS-E) Adult Immunization Status

(continued)

### Tips and Best Practices

- ✓ If a member had anaphylaxis due to any of the required vaccines, encephalitis due to Td/Tdap, or history of hepatitis B illness, submit codes in the table above to count towards the measure
- ✓ Enter administered immunizations in the Michigan Care Improvement Registry (MCIR). Information regarding MCIR can be found here: <https://mcir.org/training-resources/healthcare-provider-mcir-resources/>
- ✓ Try to complete immunizations at scheduled annual exams to reduce the number of appointments
- ✓ Offer immunizations on-site or support scheduling at accessible in-network facility during patient visit and have referrals easily accessible or standing orders
- ✓ Educate patients on the diseases that are prevented with immunizations

### Required Exclusions

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above:

- ✓ Members who use hospice services any time during the measurement year
- ✓ Members who die any time during the measurement year

---



# (BPD) Blood Pressure Control for Patients With Diabetes

## Product Lines:

- ✓ Meridian (Medicaid)
- ✓ MI Coordinated Health (Dual Eligible Plan)
- ✓ WellCare (Medicare)

## Description

Measure evaluates percentage of members 18-75 years of age with diabetes (type 1 and type 2) whose blood pressure was adequately controlled (<140/90 mm Hg) during the measurement year.

Description	Codes*
Diastolic Less Than 80	<b>CPT-CAT-II:</b> 3078F
Diastolic 80-89	<b>CPT-CAT-II:</b> 3079F
Diastolic Greater Than/Equal To 90	<b>CPT-CAT-II:</b> 3080F
Systolic Less Than 130	<b>CPT-CAT-II:</b> 3074F
Systolic 130-139	<b>CPT-CAT-II:</b> 3075F
Systolic Greater Than/Equal To 140	<b>CPT-CAT-II:</b> 3077F
Palliative Care	<b>HCPCS:</b> G9054
Hospice Care	<b>HCPCS:</b> G9473-G9479, Q5003-Q5008, Q5010, S9126, T2042-T2046, G0182 <b>CPT:</b> 99377, 99378

\*Codes subject to change

## Action

- ✓ Take member's blood pressure at each outpatient visit
- ✓ Indicate in the medical record all recorded blood pressure measurements, blood pressure results taken by the patient at home and reviewed via telehealth, or member-reported blood pressure readings

(continued)



## (BPD) Blood Pressure Control for Patients With Diabetes (continued)

### Tips and Best Practices

- ✓ If BP result is >140/90 mmHg, recheck the BP at end of the visit and document
- ✓ Ensure you are coding from the chart above for both the systolic and diastolic values
- ✓ Member-reported blood pressure readings and blood pressure readings collected via telehealth are acceptable if documented in chart with date and value. Ensure your EMR has appropriate fields to house this data for all visit types.
- ✓ Follow the American Medical Association's seven simple tips to get an accurate blood pressure reading at:  
**[www.ama-assn.org/delivering-care/hypertension/  
one-graphic-you-need-accurate-blood-pressure-reading](http://www.ama-assn.org/delivering-care/hypertension/one-graphic-you-need-accurate-blood-pressure-reading)**
- ✓ Your patient may be eligible for automatic home blood pressure monitor through their standard Over-the-Counter benefits when supplied through in-network durable medical equipment provider

### Required Exclusions

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above or medical record documentation:

- ✓ Members who use hospice services any time during the measurement year
- ✓ Members who die any time during the measurement year
- ✓ Members who receive palliative care any time during the measurement year



# (CBP) Controlling High Blood Pressure

## Product Lines:

- ✓ Meridian (Medicaid)
- ✓ MI Coordinated Health (Dual Eligible Plan)
- ✓ Ambetter from Meridian (Marketplace)
- ✓ WellCare (Medicare)

## Description

Measure evaluates the percentage of members 18-85 years of age who had a diagnosis of hypertension and whose BP was adequately controlled (<140/90 mm Hg) during the measurement year.

Description	Codes*
Diastolic Less Than 80	<b>CPT-CAT-II:</b> 3078F
Diastolic 80-89	<b>CPT-CAT-II:</b> 3079F
Diastolic Greater Than/ Equal To 90	<b>CPT-CAT-II:</b> 3080F
Systolic Less Than 130	<b>CPT-CAT-II:</b> 3074F
Systolic 130-139	<b>CPT-CAT-II:</b> 3075F
Systolic Greater Than/ Equal To 140	<b>CPT-CAT-II:</b> 3077F
Palliative Care	<b>HCPCS:</b> G9054
Hospice Care	<b>HCPCS:</b> G9473-G9479, Q5003-Q5008, Q5010, S9126, T2042-T2046, G0182 <b>CPT:</b> 99377, 99378
Members With a Diagnosis or Procedure That Indicates ESRD	<b>CPT:</b> 50220, 50225, 50230, 50234, 50236, 50340, 50370, 50543, 50545, 50546, 50548, 90935, 90937, 90945, 90947, 90997, 90999, 99512, 50240, 50360, 50365, 50380 <b>HCPCS:</b> G0257, S9339 <b>ICD-10:</b> N18.5, N18.6, Z99.2, Z94.0 <b>ICD-9:</b> 585.5, 585.6, V45.11, V42.0
Pregnancy Diagnosis	<b>ICD-10:</b> Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82, Z34.83, Z34.91, Z34.92, Z34.93

\*Codes subject to change. List does not include all codes for pregnancy, due to amount.



## (CBP) Controlling High Blood Pressure

(continued)

### Action

- ✓ Take member's blood pressure at each outpatient visit
- ✓ Indicate in the medical record all recorded blood pressure measurements, blood pressure results taken by the patient at home and reviewed via telehealth, or member-reported blood pressure readings

### Tips and Best Practices

- ✓ If BP result is  $>140/90$  mmHg, recheck the BP at end of the visit and document
- ✓ Ensure you are coding from the chart above for both the systolic and diastolic values
- ✓ Member-reported blood pressure readings and blood pressure readings collected via telehealth are acceptable if documented in chart with date and value. Ensure your EMR has appropriate fields to house this data for all visit types.
- ✓ Follow the American Medical Association's seven simple tips to get an accurate blood pressure reading at:  
**[www.ama-assn.org/delivering-care/hypertension/one-graphic-you-need-accurate-blood-pressure-reading](http://www.ama-assn.org/delivering-care/hypertension/one-graphic-you-need-accurate-blood-pressure-reading)**
- ✓ Your patient may be eligible for automatic home blood pressure monitor through their standard Over-the-Counter benefits when supplied through in-network durable medical equipment provider

### Required Exclusions

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above or medical record documentation:

- ✓ Members who use hospice services any time during the measurement year
- ✓ Members who die any time during the measurement year

(continued)



## (CBP) Controlling High Blood Pressure

(continued)

- ✓ Members who receive palliative care any time during the measurement year
- ✓ Members with evidence of end-stage renal disease (ESRD), dialysis, nephrectomy or kidney transplant any time during member's history
- ✓ Members with a diagnosis of pregnancy any time during the measurement year

.....



## (COA) Care for Older Adults

### Product Lines:

- ✓ MI Coordinated Health (Dual Eligible Plan)
- ✓ WellCare (Medicare)

### Description

Measure evaluates percentage of adults 66 years and older who had both of the following during the measurement year:

- ✓ Medication review
- ✓ Functional status assessment

Description	Codes*
Medication Review (codes must be submitted for both Medication List AND Medication Review for credit)	<b>Medication List -</b> <b>CPT-CAT-II:</b> 1159F <b>HCPCS:</b> G8427  <b>Medication Review -</b> <b>CPT-CAT-II:</b> 1160F <b>CPT:</b> 90863, 99605, 99606, 99483

(continued)



## (COA) Care for Older Adults

(continued)

Description	Codes*
Transitional Care Management Services (can be used instead of medication list/medication review codes)	<b>CPT:</b> 99495, 99496
<b>Functional Status Assessment</b>	<b>CPT:</b> 99483 <b>CPT-CAT-II:</b> 1170F <b>HCPCS:</b> G0438, G0439
<b>Hospice Care</b>	<b>HCPCS:</b> G9473–G9479, Q5003–Q5008, Q5010, S9126, T2042–T2046, G0182 <b>CPT:</b> 99377, 99378

\*Codes subject to change

### Action

#### ✓ Medication review

- Perform an annual review of the patient's medications, including prescription medications, over-the-counter medications and herbal or supplemental therapies. This must be completed by a prescribing provider or a clinical pharmacist

#### ✓ Functional status assessment

- Assess the patient's ability to perform daily tasks and identify any functional decline
- This can be documented through assessing ADLS, IADLS, or using a standardized functional status assessment tool

### Tips and Best Practices

- ✓ If member is not taking any medication, document this in the record to count for medication review

(continued)



## (COA) Care for Older Adults

*(continued)*

- ✓ No specific setting or provider specialty is required for functional status assessment. Telephone visit, e-visit or virtual check-in meet criteria
- ✓ Complete medication review and functional status assessment in the same visit when able. Annual wellness visits are a great opportunity for this

### **Required Exclusions**

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above or medical record documentation:

- ✓ Members who use hospice services any time during the measurement year
- ✓ Members who die any time during the measurement year

.....



## (COL-E) Colorectal Cancer Screening

### **Product Lines:**

- ✓ Meridian (Medicaid)
- ✓ MI Coordinated Health (Dual Eligible Plan)
- ✓ Ambetter from Meridian (Marketplace)
- ✓ WellCare (Medicare)

### **Description**

Measure evaluates the percentage of members 45-75 years of age who had appropriate screening for colorectal cancer.

*(continued)*



## (COL-E) Colorectal Cancer Screening

(continued)

Description	Codes*
Colonoscopy	<b>CPT:</b> 44388-44392, 44394, 44401-44408, 45378-45382, 45384-45386, 45388-45393, 45398 <b>HCPCS:</b> G0105, G0121
CT Colonography	<b>CPT:</b> 74261-74263
sDNA Fit Lab Test (Cologuard)	<b>CPT:</b> 81528, 0464U <b>LOINC:</b> 77353-1, 77354-9
Flexible Sigmoidoscopy	<b>CPT:</b> 45330-45335, 45337, 45338, 45340, 45341, 45342, 45346, 45347, 45349, 45350 <b>HCPCS:</b> G0104
FOBT Lab Test	<b>CPT:</b> 82270, 82274 <b>HCPCS:</b> G0328
Palliative Care	<b>HCPCS:</b> G9054
Hospice Care	<b>HCPCS:</b> G9473-G9479, Q5003-Q5008, Q5010, S9126, T2042-T2046, G0182 <b>CPT:</b> 99377, 99378
Colorectal Cancer	<b>ICD-10:</b> C18.0, C18.1, C18.2, C18.3, C18.4, C18.5 C18.6, C18.7, C18.8, C18.9, C19, C20, C21.2, C21.8, C78.5, Z85.038, Z85.048
Total Colectomy	<b>CPT:</b> 44150-44153, 44155-44158, 44210-44212

\*Codes subject to change

### Action

Inform your patient that while a colonoscopy needs to be done the least frequently, there are multiple options for colorectal cancer screenings:

- ✓ Colonoscopy: during the measurement year or nine years prior
- ✓ Stool DNA (sDNA) with FIT Test or Cologuard®: during the measurement year or two years prior
- ✓ Fecal occult blood test (FOBT): during the measurement year
- ✓ Flexible sigmoidoscopy: during the measurement year or four years prior
- ✓ CT colonography: during the measurement year or four years prior



## (COL-E) Colorectal Cancer Screening

(continued)

### Tips and Best Practices

- ✓ Member-reported screenings are acceptable if documented in chart as part of member's history. It must be clear the screening happened in timeframe, but an exact date and result is not needed (for example: "colonoscopy done about 5 years ago" or "colonoscopy within last 3 years" are both acceptable)
- ✓ Support colonoscopy scheduling at accessible in-network facility during patient visit and have referrals easily accessible or standing orders
- ✓ Reach out to your Quality Improvement or Provider Network Management Representative to find out if your patient received a FIT home test kit from the health plan

### Required Exclusions

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above or medical record documentation:

- ✓ Members who use hospice services any time during the measurement year
- ✓ Members who die any time during the measurement year
- ✓ Members who receive palliative care any time during the measurement year
- ✓ Members who had colorectal cancer any time during the member's history
- ✓ Members who had a total colectomy any time during the member's history



## (EED) Eye Exam for Patients With Diabetes

### Product Lines:

- ✓ Meridian (Medicaid)
- ✓ MI Coordinated Health (Dual Eligible Plan)
- ✓ Ambetter from Meridian (Marketplace)
- ✓ Wellcare (Medicare)

### Description

Measure evaluates percentage of members 18-75 years of age with diabetes (type 1 and type 2) who had one of the following completed by an eye care professional (optometrist or ophthalmologist):

- a retinal or dilated eye exam during the measurement year
- a retinal or dilated eye exam with negative findings (negative for retinopathy) in the year prior

Description	Codes*
<b>Eye Exam with Evidence of Retinopathy</b>	<b>CPT:</b> 2022F, 2024F, 2026F
<b>Eye Exam Without Evidence of Retinopathy</b>	<b>CPT:</b> 2023F, 2025F, 2033F
<b>Autonomous Eye Exam</b>	<b>CPT:</b> 92229 <b>CPT-LOINC:</b> 105914-6 with a result
<b>Retinal Eye Exams (must be billed by eye care professional and only counts for measurement year; add a code from Diabetes Mellitus Without Complications set below to last for two years)</b>	<b>CPT:</b> 92002, 92004, 92012, 92014, 92018, 92019, 92134, 92137, 92201, 92202, 92230, 92235, 92250, 99203-99205, 99213-99215, 99242-99245 <b>HCPCS:</b> S0620, S0621, S3000
<b>Diabetes Mellitus Without Complications</b>	<b>ICD-10:</b> E10.9, E11.9, E13.9
<b>Retinal Imaging With Interpretation and Reporting by Qualified Reading Center</b>	<b>CPT:</b> 92227, 92228
<b>Diabetic Retinal Screening Negative in Prior Year</b>	<b>CPT:</b> 3072F



## (EED) Eye Exam for Patients With Diabetes

(continued)

Description	Codes*
<b>Unilateral Eye Enucleation With a Bilateral Modifier (must add CPT Modifier: 50)</b>	<b>CPT:</b> 65091, 65093, 65101, 65103, 65105, 65110, 65112, 65114
<b>Bilateral Absence of Eyes</b>	<b>SNOMED:</b> 15665641000119103
<b>Palliative Care</b>	<b>HCPCS:</b> G9054
<b>Hospice Care</b>	<b>HCPCS:</b> G9473–G9479, Q5003–Q5008, Q5010, S9126, T2042–T2046, G0182 <b>CPT:</b> 99377, 99378

\*Codes subject to change

### Action

- ✓ Refer your diabetic members to an eye care professional annually to complete a retinal or dilated eye exam or complete in office and send exam to ophthalmologist or optometrist for review and sign off
- ✓ If patient tests negative for retinopathy, eye exam can be performed every other year

### Tips and Best Practices

- ✓ Conduct outreach to diabetic population about yearly Dilated Retinal Exams
- ✓ Remind patients concerned with costs of Dilated Retinal Exams that it's a covered medical benefit
- ✓ Support exam scheduling at accessible in-network facility during patient visit and have referrals easily accessible or standing orders
- ✓ Reach out to your Quality or Provider Network Management Representative to find out about retinal camera rental or if your patient is eligible for in-home retinal screening

(continued)



## (EED) Eye Exam for Patients With Diabetes

(continued)

### Required Exclusions

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above or medical record documentation:

- ✓ Members who had a bilateral eye enucleation any time during the member's history
- ✓ Members who have bilateral absence of eyes any time during the member's history
- ✓ Members who die any time during the measurement year
- ✓ Members who receive palliative care any time during the measurement year
- ✓ Members who use hospice services any time during the measurement year

---



## (FMC) Follow-Up After Emergency Department Visit for People With Multiple High-Risk Chronic Conditions

### Product Lines:

- ✓ WellCare (Medicare)
- ✓ MI Coordinated Health (Dual Eligible Plan)

### Description

Measure evaluates the percentage of emergency department (ED) visits for persons 18 years of age and older who have multiple high-risk chronic conditions who had a follow-up service within 7 days of the ED visit.



## (FMC) Follow-Up After Emergency Department Visit for People With Multiple High-Risk Chronic Conditions *(continued)*

Patients must have had two or more of the following chronic conditions documented on two different dates of services to be included:

- COPD, asthma or unspecified bronchitis
- Alzheimer's disease and related disorders
- Chronic kidney disease
- Depression
- Heart failure
- Acute myocardial infarction
- Atrial fibrillation
- Stroke and transient ischemic attack

Description	Codes*
<b>Outpatient Visit, Telephone Visit, E-visit, or Virtual Check-In</b>	<b>CPT:</b> 98000-98016, 98966-98968, 98970-98972, 98980, 98981, 99202-99205, 99211-99215, 99242-99245, 99341, 99342, 99344, 99345, 99347-99350, 99381-99387, 99391-99397, 99401-99404, 99411, 99412, 99421-99423, 99492, 99441-99443, 99455- 99458, 99483 <b>HCPCS:</b> G0071, G0402, G0438, G0439, G0463, G2010, G2012, G2250, G2251, G2252, T1015
<b>Transitional Care Management</b>	<b>CPT:</b> 99495, 99496
<b>Case Management Visits</b>	<b>CPT:</b> 99366 <b>HCPCS:</b> T1016, T1017, T2022, T2023
<b>Complex Care Management Services</b>	<b>CPT:</b> 99439, 99487, 99489, 99490, 99491 <b>HCPCS:</b> G0506



# (FMC) Follow-Up After Emergency Department Visit for People With Multiple High-Risk Chronic Conditions (continued)

Description	Codes*
<b>BH Outpatient Visit</b>	<b>CPT:</b> 98000-98007, 98960-98962, 99078, 99202-99205, 99211-99215, 99242-99245, 99341-99345, 99347-99350, 99381-99387, 99391-99397, 99401-99404, 99411, 99412, 99483, 99492-99494, 99510 <b>HCPCS:</b> G0155, G0176, G0177, G0409, G0463, G0512, G0560, H0002, H0004, H0031, H0034, H0036, H0037, H0039, H0040, H2000, H2010, H2011, H2013-H2020, T1015
<b>Partial Hospitalization or Intensive Outpatient</b>	<b>HCPCS:</b> G0410, G0411, H0035, H2001, H2012, S0201, S9480, S9484, S9485
<b>Electroconvulsive Therapy</b>	<b>CPT:</b> 90870
<b>Substance Use Disorder Service</b>	<b>CPT:</b> 99408, 99409 HCPCS: G0396, G0397, G0443, H0001, H0005, H0007, H0015, H0016, H0022, H0047, H0050, H2034, H2036, T1006, T1012
<b>Substance Abuse Counseling and Surveillance</b>	<b>ICD10CM:</b> Z71.41, Z71.51
<b>Hospice Care</b>	<b>HCPCS:</b> G9473-G9479, Q5003-Q5008, Q5010, S9126, T2042-T2046, G0182 <b>CPT:</b> 99377, 99378

\*Codes subject to change

## Action

✓ Complete one of the following within seven days of ED visit:

- Outpatient visit, telephone visit, e-visit, or virtual check-in
- Transitional care management services



## (FMC) Follow-Up After Emergency Department Visit for People With Multiple High-Risk Chronic Conditions *(continued)*

- Case management visit
- Complex Care Management service
- Outpatient or telehealth behavioral health visit
- Intensive outpatient encounter or partial hospitalization
- Community mental health center visit
- Electroconvulsive therapy
- Substance use disorder service
- Substance use disorder counseling and surveillance

### Tips and Best Practices

- ✓ Implement processes with hospitals to facilitate timely sharing of ED disposition information
- ✓ Educate and encourage patients to notify their primary care physician following an ED visit for additional guidance
- ✓ Schedule the seven-day follow-up visit within five days to allow flexibility in rescheduling. Involve the patient or patient's caregiver regarding the follow-up plan
- ✓ Follow-up visit can occur on same day of ED visit
- ✓ There is not a specific provider type requirement for this measure

### Required Exclusions

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above or medical record documentation:

- ✓ Members who use hospice services any time during the measurement year
- ✓ Members who die anytime during the measurement year



# (GSD) Glycemic Status Assessment for Patients With Diabetes

## Product Lines:

- ✓ Meridian (Medicaid)
- ✓ MI Coordinated Health (Dual Eligible Plan)
- ✓ Ambetter from Meridian (Marketplace)
- ✓ WellCare (Medicare)

## Description

Measure evaluates percentage of members 18-75 years of age with diabetes (type 1 and type 2) whose most recent glycemic status was at the following levels during the measurement year:

- ✓ HbA1c or glucose management indicator control (<8.0%)
- ✓ HbA1c or glucose management indicator poor control (>9.0%)

Description	Codes*
HbA1c Level Less Than 7	<b>CPT-CAT-II:</b> 3044 F
HbA1c Level Greater Than/ Equal to 7 and Less Than 8	<b>CPT-CAT-II:</b> 3051F
HbA1c Level Greater Than/ Equal to 8 and Less Than/ Equal to 9	<b>CPT-CAT-II:</b> 3052F
HbA1C Greater Than 9	<b>CPT-CAT-II:</b> 3046F
Palliative Care	<b>HCPCS:</b> G9054
Hospice Care	<b>HCPCS:</b> G9473–G9479, Q5003–Q5008, Q5010, S9126, T2042–T2046, G0182 <b>CPT:</b> 99377, 99378

\*Codes subject to change

## Action

- ✓ Perform at least one HbA1c test annually or monitor your patient with a Continuous Glucose Monitoring system
- ✓ For results >8%, repeat the test later in the measurement year



## (GSD) Glycemic Status Assessment for Patients With Diabetes *(continued)*

### Tips and Best Practices

- ✓ If there are multiple glycemic status assessments on the same date of service, use the lowest result
- ✓ Member-reported HbA1c results are acceptable if documented in chart with test date and value. Ensure your EMR has appropriate fields to house this data for all visit types
- ✓ Discuss the importance of HbA1c/blood glucose control in patients with diabetes and long-term effects of elevated blood glucose
- ✓ Offer referrals to Diabetes Education, Endocrinologist, and/or Dietician
- ✓ Offer lab testing on-site or support lab scheduling at accessible in-network facility during patient visit and have referrals easily accessible or standing orders
- ✓ Reach out to your Quality or Provider Network Management Representative to find out if your patient received a HbA1c home test kit from the health plan
- ✓ For patients taking diabetes medication, encourage adherence by providing 90-day prescriptions. Your patient may be eligible for prescription delivery by mail through Express Scripts
- ✓ Your patient may be eligible for diabetes education services through an approved in network provider

### Required Exclusions

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above or medical record documentation:

- ✓ Members who use hospice services any time during the measurement year
- ✓ Members who die any time during the measurement year
- ✓ Members who receive palliative care any time during the measurement year

.....

*(continued)*



# (KED) Kidney Health Evaluation for Patients With Diabetes

## Product Lines:

- ✓ Meridian (Medicaid)
- ✓ MI Coordinated Health (Dual Eligible Plan)
- ✓ Ambetter from Meridian (Marketplace)
- ✓ WellCare (Medicare)

## Description

The percentage of members 18–85 years of age with diabetes (type 1 and type 2) who received a kidney health evaluation, defined by an estimated glomerular filtration rate (eGFR) and a urine albumin-creatinine ratio (uACR), during the measurement year.

Description	Codes*
Estimated Glomerular Filtration Rate (eGFR)	<b>CPT:</b> 80047, 80048, 80050, 80053, 80069, 82565
Urine Creatinine Lab Test	<b>CPT:</b> 82570
Quantitative Urine Albumin Lab Test	<b>CPT:</b> 82043
Urine Albumin Creatinine Ratio Lab Test	<b>LOINC:</b> 13705-9, 14958-3, 14959-1, 30000-4, 44292-1, 59159-4, 76401-9, 77253-3, 77254-1, 89998-9, 9318-7
ESRD Diagnosis	<b>ICD:</b> N18.5, N18.6
Dialysis Procedure	<b>CPT:</b> 90935, 90937, 90945, 90947, 90997, 90999, 99512 <b>HCPCS:</b> G0257, S9339
Palliative Care	<b>HCPCS:</b> G9054
Hospice Care	<b>HCPCS:</b> G9473–G9479, Q5003–Q5008, Q5010, S9126, T2042–T2046, G0182 <b>CPT:</b> 99377, 99378

\*Codes subject to change

(continued)



# (KED) Kidney Health Evaluation for Patients With Diabetes *(continued)*

## Action

- ✓ Perform both eGFR and uACR testing annually for patients with diabetes. UACR testing can be either of the following:
  - Urine Albumin Creatinine Ratio Lab Test
  - Both a quantitative urine albumin lab test and a urine creatinine test with service dates four days or less apart

## Tips and Best Practices

- ✓ Member-reported results are acceptable if documented in chart with test date and value. Ensure your EMR has appropriate fields to house this data for all visit types.
- ✓ This is not a measure where the results of the tests will affect compliance. The intent is that the services were completed
- ✓ Encourage members to have annual testing for appropriate identification, staging, monitoring and treatment of kidney disease
- ✓ Offer lab testing on-site or support lab scheduling at accessible in-network facility during patient visit and have referrals easily accessible or standing orders
- ✓ Your patient may be eligible for diabetes education services through an approved in network provider
- ✓ Reach out to your Quality or Provider Network Management Representative to find out if your patient received an eGFR and uACR home test kit from the health plan

## Required Exclusions

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above or medical record documentation:

- ✓ Members who use hospice services any time during the measurement year
- ✓ Members who die any time during the measurement year

*(continued)*



## (KED) Kidney Health Evaluation for Patients With Diabetes (continued)

- ✓ Members who receive palliative care any time during the measurement year
- ✓ Members who have evidence of ESRD or dialysis any time during the member's history

.....



## (PCE) Pharmacotherapy Management of COPD Exacerbation

### **Product Lines:**

- ✓ Meridian (Medicaid)
- ✓ MI Coordinated Health (Dual Eligible Plan)
- ✓ WellCare (Medicare)

### **Description**

Measure evaluates the percentage of COPD exacerbations for members 40 years of age and older who had an acute inpatient discharge or emergency department visit on or between January 1 to November 30 of the measurement year and who were dispensed one of the following appropriate medications or had evidence of an active prescription:

1. A systemic corticosteroid within 14 days of inpatient discharge or ED visit
2. A bronchodilator within 30 days of inpatient discharge or ED visit



# (PCE) Pharmacotherapy Management of COPD Exacerbation (continued)

Description	Codes*
Hospice Care	<b>HCPCS:</b> G9473-G9479, Q5003-Q5008, Q5010, S9126, T2042-T2046, G0182 CPT: 99377, 99378

## Systemic Corticosteroid Medications

Description	Prescriptions	
Glucocorticoids	<ul style="list-style-type: none"><li>Cortisone</li><li>Dexamethasone</li><li>Hydrocortisone</li></ul>	<ul style="list-style-type: none"><li>Methylprednisolone</li><li>Prednisolone</li><li>Prednisone</li></ul>

## Bronchodilator Medications

Description	Prescriptions	
Anticholinergic agents	<ul style="list-style-type: none"><li>Aclidinium bromide</li><li>Ipratropium</li></ul>	<ul style="list-style-type: none"><li>Tiotropium</li><li>Umeclidinium</li></ul>
Beta 2-agonists	<ul style="list-style-type: none"><li>Albuterol</li><li>Arformoterol</li><li>Formoterol</li><li>Indacaterol</li></ul>	<ul style="list-style-type: none"><li>Levalbuterol</li><li>Metaproterenol</li><li>Oladaterol</li><li>Salmeterol</li></ul>
Bronchodilator combinations	<ul style="list-style-type: none"><li>Albuterol-ipratropium</li><li>Budesonide-formoterol</li><li>Fluticasone-salmeterol</li><li>Fluticasone-vilanterol</li><li>Fluticasone furoate-umeclidinium-vilanterol</li><li>Formoterol-aclidinium</li><li>Formoterol-glycopyrrolate</li></ul>	<ul style="list-style-type: none"><li>Formoterol-mometasone</li><li>Glycopyrrolate-indacaterol</li><li>Oladaterol-tiotropium</li><li>Umeclidinium-vilanterol</li></ul>

## Action

- ✓ Schedule a follow-up appointment after discharge and confirm that the patient has been prescribed the appropriate medication and had it filled

(continued)



## (PCE) Pharmacotherapy Management of COPD Exacerbation (continued)

### Tips and Best Practices

- ✓ Reconcile patients' medications with those prescribed at discharge when you receive the discharge summary
- ✓ Ask the patient if they have any barriers that prevent them from filling their prescriptions
- ✓ Provide a COPD action plan for the patient, including daily medications, trigger avoidance, and what to do when flare-ups occur. This resource from the American Lung Association can be referenced: <https://www.lung.org/lung-health-diseases/lung-disease-lookup/copd/living-with-copd/copd-management-tools>

### Required Exclusions

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above or medical record documentation:

- ✓ Members who use hospice services any time during the measurement year
- ✓ Members who die any time during the measurement year

---



## (PCR) Plan All-Cause Readmissions

### Product Lines:

- ✓ Meridian (Medicaid)
- ✓ MI Coordinated Health (Dual Eligible Plan)
- ✓ Ambetter from Meridian (Marketplace)
- ✓ WellCare (Medicare)



## (PCR) Plan All-Cause Readmissions (continued)

### Description

For members 18 years of age and older, the number of acute inpatient and observation stays during the measurement year that were followed by an unplanned acute readmission for any diagnosis with 30 days and the predicted probability of an acute readmission.

Description	Codes*
Hospice Care	<b>HCPCS:</b> G9473-G9479, Q5003-Q5008, Q5010, S9126, T2042-T2046, G0182 <b>CPT:</b> 99377, 99378

\*Codes subject to change

### Action

- ✓ Coordinate care from the hospital to home and ensure a follow-up visit with the primary care physician to help your patients avoid a readmission

### Tips and Best Practices

- ✓ Implement a post-discharge process to track, monitor and follow up with patients
- ✓ Provide transitional care management for patients who are at high-risk for readmissions
- ✓ Keep a few open appointments available so patients who are discharged from the hospital can be seen within 7 days of discharge
- ✓ Discuss the discharge summary with your patients and ask if they understand the instructions and filled new prescriptions

### Required Exclusions

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above or medical record documentation:

- ✓ Members who use hospice services any time during the measurement year
- ✓ Members who die any time during the measurement year



# (SSD) Diabetes Screening for People with Schizophrenia or Bipolar Disorder Who are Using Antipsychotic Medication

## Product Lines:

- ✓ Meridian (Medicaid)

## Description

Measure evaluates percentage of members 18-64 years of age with schizophrenia, schizoaffective disorder or bipolar disorder who were dispensed an antipsychotic medication and had a diabetes screening test during the measurement year.

DESCRIPTION	CODES*
HbA1c Lab Tests	<b>CPT:</b> 83036, 83037 <b>CPT-CAT-II:</b> 3046F, 3051F, 3052F, 3044F
Glucose Lab Tests	<b>CPT:</b> 80047, 80048, 80050, 80053, 80069, 82947, 82951, 82950
Hospice Care	<b>HCPCS:</b> G9473-G9479, Q5003-Q5008, Q5010, S9126, T2042-T2046, G0182 <b>CPT:</b> 99377, 99378
Diabetes	<b>ICD-10:</b> E10.8, E10.9, E11.8, E11.9 **list does not include all codes for diabetes, due to amount

\*Codes subject to change

## Action

- ✓ Ensure patients who have one or more antipsychotic prescriptions have HbA1c testing or glucose testing completed in the measurement year

## Tips and Best Practices

- ✓ Educate patients on risks of hyperglycemia associated with antipsychotic medication treatment
- ✓ Address concerns regarding side effects and how to manage them
- ✓ Offer lab testing on-site or support lab scheduling at accessible in-network facility during patient visit and have referrals easily accessible or standing orders



## (SSD) Diabetes Screening for People with Schizophrenia or Bipolar Disorder Who are Using Antipsychotic Medication *(continued)*

### Required Exclusions

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above or medical record documentation:

- ✓ Members who use hospice services any time during the measurement year
- ✓ Members who die any time during the measurement year
- ✓ Members with diabetes, identified by one of the following:
  - At least two diagnoses of diabetes during the measurement year or the year prior
  - At least one of diagnosis of diabetes and dispensed insulin or hypoglycemics/antihyperglycemics at least once during the measurement year or year prior
- ✓ Members who had no antipsychotic medications dispensed during the measurement year

.....



## (TRC) Transitions of Care

### Product Lines:

- ✓ MI Coordinated Health (Dual Eligible Plan)
- ✓ WellCare (Medicare)

### Description

Measure evaluates percentage of discharges for members 18 years of age and older who had each of the following:

- ✓ Notification of Inpatient Admission: Documentation of receipt of notification of inpatient admission on the day of admission through 2 days after the admission
- ✓ Receipt of Discharge Information: Documentation of receipt of discharge information on the day of discharge through 2 days after the discharge
- ✓ Patient Engagement After Inpatient Discharge: Documentation of patient engagement provided within 30 days after discharge
- ✓ Medication Reconciliation Post-Discharge: Documentation of medication reconciliation conducted by a prescribing practitioner, clinical pharmacist, physician assistant or registered nurse on the date of discharge through 30 days after discharge

DESCRIPTION	CODES*
<b>Medication Reconciliation</b>	<b>CPT:</b> 99483, 99495, 99496, 99605, 99606 <b>CPT-CAT-II:</b> 1111F
<b>Outpatient and Telehealth (for Patient Engagement)</b>	<b>CPT:</b> 98000-98016, 98966-98968, 98970-98972, 98980, 98981, 99202-99205, 99211-99215, 99242-99245, 99341, 99342, 99344, 99345, 99347-99350, 99381-99387, 99391-99397, 99401-99404, 99411, 99412, 99421-99423, 99429, 99441-99443, 99455-99458, 99483 <b>HCPCS:</b> G0071, G0402, G0438, G0439, G0463, G2010, G2012, G2250, G2251, G2252, T1015

(continued)



## (TRC) Transitions of Care (continued)

DESCRIPTION	CODES*
<b>Transitional Care Management Services (for Patient Engagement)</b>	<b>CPT:</b> 99495, 99496
<b>Hospice Care</b>	<b>HCPCS:</b> G9473-G9479, Q5003-Q5008, Q5010, S9126, T2042-T2046, G0182 <b>CPT:</b> 99377, 99378

\*Codes subject to change

### Action

- ✓ Document in the member's medical record the date the PCP was notified of the member's in-patient hospital admission
- ✓ Obtain discharge summaries from inpatient facilities within 3 days of discharge
- ✓ Reach out to member immediately after you receive notification that they have been discharged to set up a follow-up visit within 30 days. This can be an office visit, home visit, phone call or telehealth visit
- ✓ During follow-up visit, review all of member's current medications compared to medications prior to admission

### Tips and Best Practices

- ✓ Investigate and implement a process to receive auto alerts when a member is admitted to or discharged from an inpatient facility
- ✓ Date and time stamp the receipt of notification admission in the member's medical record
- ✓ Date and time stamp the receipt of the discharge summary in the member's medical record
- ✓ Schedule follow-up appointment with primary care provider within the first seven days of discharge, to allow time for rescheduling if needed
- ✓ Reconcile the patient discharge medications to his or her outpatient medications at follow-up visit



## (TRC) Transitions of Care *(continued)*

### Required Exclusions

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above or medical record documentation:

- ✓ Members who use hospice services any time during the measurement year
- ✓ Members who die any time during the measurement year

.....

# Women's Health



## (BCS-E) Breast Cancer Screening

### Product Lines:

- ✓ Meridian (Medicaid)
- ✓ MI Coordinated Health (Dual Eligible Plan)
- ✓ Ambetter from Meridian (Marketplace)
- ✓ WellCare (Medicare)

### Description

Measure evaluates the percentage of members 40-74 years of age who were recommended for routine breast cancer screening and had a mammogram to screen for breast cancer.

Description	Codes*
Mammography	<b>CPT:</b> 77061-77063, 77065-77067
History of Bilateral Mastectomy	<b>ICD-10:</b> Z90.13
Bilateral Mastectomy	<b>ICD10PCS:</b> OHTVOZZ
Unilateral Mastectomy with Bilateral Modifier <b>(add CPT Modifier code 50 if bilateral done in same procedure)</b>	<b>CPT:</b> 19180, 19200, 19220, 19240, 19303, 19304, 19305, 19306, 19307
Absence of Left Breast	<b>ICD-10:</b> Z90.12
Absence of Right Breast	<b>ICD-10:</b> Z90.11
Gender-Affirming Chest Surgery	<b>CPT:</b> 19318
Gender Dysphoria	<b>ICD-10:</b> F64.1, F64.2, F64.8, F64.9, Z87.890

(continued)



## (BCS-E) Breast Cancer Screening (continued)

Palliative Care	<b>HCPCS:</b> G9054, M1017
Hospice Care	<b>HCPCS:</b> G9473-G9479, Q5003-Q5008, Q5010, S9126, T2042-T2046, G0182 <b>CPT:</b> 99377, 99378

\*Codes subject to change

### Action

- ✓ Make sure your patient completes a mammogram at least every two years

### Tips and Best Practices

- ✓ Member-reported screenings are acceptable if documented in chart with date. It must be clear the screening happened in timeframe, but an exact date and result is not needed (for example: “mammogram done about 4 months ago” or “mammogram within last year” are both acceptable). Ensure your EMR has appropriate fields to house this data for all visit types.
- ✓ Support mammogram scheduling at accessible in-network facility during patient visit and have referrals easily accessible or standing orders
- ✓ All types and methods of mammograms are acceptable, including screening, diagnostic, film, digital or digital tomosynthesis

### Required Exclusions

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above or medical record documentation:

- ✓ Members who use hospice services any time during the measurement year
- ✓ Members who die any time during the measurement year
- ✓ Members who received palliative care during the measurement year



## (BCS-E) Breast Cancer Screening (continued)

- ✓ Members who had a bilateral mastectomy or both right and left unilateral mastectomies any time during the member's history
- ✓ Members who had gender-affirming chest surgery with a diagnosis of gender dysphoria any time during the member's history

.....



## (CCS-E) Cervical Cancer Screening

### Product Lines:

- ✓ Meridian (Medicaid)
- ✓ Ambetter from Meridian (Marketplace)

### Description

This measure demonstrates the percentage of members 21-64 years of age who were recommended for routine cervical cancer screening and were screened based on the following criteria:

- ✓ Members 21-64 years of age who had cervical cytology performed within the last 3 years.
- ✓ Members 30-64 years of age who had cervical high-risk human papillomavirus (hrHPV) testing performed within the last 5 years.
- ✓ Members 30-64 years of age who had cervical cytology/high risk human papillomavirus (hrHPV) co-testing within the last 5 years.

Description	Codes*
<b>Cervical Cytology Lab Test (21-64)</b>	<b>CPT:</b> 88141-88143, 88147, 88148, 88150, 88152, 88153, 88164-88167, 88174, 88175 <b>HCPCS:</b> G0123, G0124, G0141, G0143, G0144, G0145, G0147, G0148, P3000, P3001



## (CCS-E) Cervical Cancer Screening (continued)

HPV Tests (30-64)	<b>CPT:</b> 87624, 87625, 87626, 0502U <b>HCPCS:</b> G0476
Hysterectomy with No Residual Cervix	<b>CPT:</b> 57530, 57531, 57540, 57545, 57550, 57555, 57556, 58150, 58152, 58200, 58210, 58240, 58260, 58262, 58263, 58267, 58270, 58275, 58280, 58285, 58290–58294, 58548, 58550, 58552–58554, 58570–58573, 58575, 58951, 58953, 58954, 58956, 59135
Absence of Cervix Diagnosis	<b>ICD-10:</b> Q51.5, Z90.710, Z90.712
Members with Sex Assigned at Birth of Male	<b>LOINC:</b> 76689-9 and LA2-8
Palliative Care	<b>HCPCS:</b> G9054
Hospice Care	<b>HCPCS:</b> G9473-G9479, Q5003-Q5008, Q5010, S9126, T2042-T2046, G0182 <b>CPT:</b> 99377, 99378

\*Codes subject to change

### Action

- ✓ Make sure your patient completes a cervical cytology every 3 years, or HPV screening every 5 years if over 30 years of age

### Tips and Best Practices

- ✓ Member-reported screenings are acceptable if documented in chart with date and results. It must be clear the screening happened in timeframe, but an exact date is not needed (for example: “pap smear done about 4 months ago, WNL” or “HPV negative 2023” are both acceptable). Ensure your EMR has appropriate fields to house this data for all visit types.
- ✓ Provide cervical cancer screening reminder at the time of appointment scheduling so the patient is aware and prepared for the exam during the visit
- ✓ Offer cervical cancer screening in office or support scheduling with accessible in-network provider during visits

(continued)



## (CCS-E) Cervical Cancer Screening *(continued)*

- ✓ Offer support to patients that are fearful of the exam and be mindful of barriers related to trauma history

### Required Exclusions

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above or medical record documentation:

- ✓ Members who use hospice services any time during the measurement year
- ✓ Members who die any time during the measurement year
- ✓ Members who received palliative care during the measurement year
- ✓ Members who had a hysterectomy with no residual cervix, cervical agenesis, or acquired absence of cervix any time during the member's history
- ✓ Members with Sex Assigned at Birth (LOINC code 76689-9) of Male (LOINC code LA2-8) at any time in the member's history

---



## (CHL) Chlamydia Screening

### Product Lines:

- ✓ Meridian (Medicaid)
- ✓ Ambetter from Meridian (Marketplace)

### Description

Measure evaluates the percentage of members 16-24 years of age who were recommended for routine chlamydia screening, were identified as sexually active and had at least one test for chlamydia during the measurement year.



## (CHL) Chlamydia Screening (continued)

Description	Codes*
Chlamydia Tests	<b>CPT:</b> 87110, 87270, 87320, 87490-87492, 87810
Hospice Care	<b>HCPGS:</b> G9473-G9479, Q5003-Q5008, Q5010, S9126, T2042-T2046, G0182 <b>CPT:</b> 99377, 99378

\*Codes subject to change

### Action

- ✓ Perform chlamydia screening each year on every 16 to 24-year-old female identified as being sexually active

### Tips and Best Practices

- ✓ Chlamydia screening can be completed via urine specimen, clinician-collected vaginal swab or self-collected vaginal swab. Providing options to patient may ease fear
- ✓ Include chlamydia screening as a part of routine clinical preventive care, walk-in visits, pregnancy testing, and emergency contraception counseling visits
- ✓ Member-reported screenings are acceptable if documented in chart with date. It must be clear the screening happened in timeframe, but an exact date and result is not needed (for example: “chlamydia screening done about 2 months ago” is acceptable). Ensure your EMR has appropriate fields to house this data for all visit types
- ✓ Meet with teens and young adults separately from their parents to allow open conversation
- ✓ Offer lab testing on-site or support lab scheduling at accessible in-network facility during patient visit and have referrals easily accessible or standing orders
- ✓ Reach out to your Quality or Provider Network Management Representative to find out if your patient received a chlamydia home test kit from the health plan

(continued)



## (CHL) Chlamydia Screening (continued)

### Required Exclusions

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above or medical record documentation:

- ✓ Members who use hospice services any time during the measurement year
- ✓ Members who die any time during the measurement year
- ✓ Members with Sex Assigned at Birth (LOINC CODE 76689-9) of Male (LOINC code LA2-8) at any time in the member's history

---



## (OMW) Osteoporosis Management in Women Who Had a Fracture

### Product Lines:

- ✓ MI Coordinated Health (Dual Eligible Plan)
- ✓ WellCare (Medicare)

### Description

Measure evaluates the percentage of women 67-85 years of age who suffered a fracture and who had either a bone mineral density (BMD) test or prescription for a drug to treat osteoporosis in the six months after the fracture.

Description	Codes*
Bone Mineral Density Tests	<b>CPT:</b> 76977, 77078, 77080, 77081, 77085, 77086
Osteoporosis Medication Therapy	<b>HCPCS:</b> J0897, J1740, J3110, J3111, J3489
Long-Acting Osteoporosis Medications during Inpatient Stay	<b>HCPCS:</b> J0897, J1740, J3489
Palliative Care	<b>HCPCS:</b> G9054

(continued)



## (OMW) Osteoporosis Management in Women Who Had a Fracture (continued)

Hospice Care	<b>HCPCS:</b> G9473-G9479, Q5003-Q5008, Q5010, S9126, T2042-T2046, G0182 <b>CPT:</b> 99377, 99378
--------------	--

\*Codes subject to change

Osteoporosis Medications		
Description	Prescription	
Bisphosphonates	• Alendronate • Ibandronate	• Alendronate-cholecalciferol • Risedronate • Zoledronic acid
Other agents	• Abaloparatide • Teriparatide	• Romosozumab • Denosumab • Raloxifene

### Action

- ✓ Make sure that all women ages 65-75 receive a routine osteoporosis screening with a BMD test. For women who have suffered a fracture, complete a BMD test or fill a medication for osteoporosis treatment within six months.

### Tips and Best Practices

- ✓ Appropriate testing or treatment for osteoporosis after the fracture is defined by any of the following criteria:
  - A BMD test on the date of fracture or in the 6-month period after fracture
  - Osteoporosis therapy on the date of fracture or in the 6-month period after the fracture
  - A dispensed prescription to treat osteoporosis on the date of fracture or in the 6-month period after the fracture
- ✓ Support BMD scheduling with accessible in-network provider during patient visit
- ✓ Reach out to your Quality or Provider Network Management Representative to find out if your patient is eligible for an in-home BMD



## (OMW) Osteoporosis Management in Women Who Had a Fracture (continued)

### Required Exclusions

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above or medical record documentation:

- ✓ Members who use hospice services any time during the measurement year
- ✓ Members who die any time during the measurement year
- ✓ Members who received palliative care during the measurement year

---



## (OSW) Osteoporosis Screening in Older Women

### Product Lines:

- ✓ MI Coordinated Health (Dual Eligible Plan)
- ✓ WellCare (Medicare)

### Description

The percentage of women 65-75 years of age who received osteoporosis screening.

Description	Codes*
Osteoporosis Screening Tests	<b>CPT:</b> 76977, 77078, 77080, 77081, 77085
Palliative Care	<b>HCPCS:</b> G9054
Hospice Care	<b>HCPCS:</b> G9473–G9479, Q5003–Q5008, Q5010, S9126, T2042–T2046, 99377, 99378
Osteoporosis Medication Therapy	<b>HCPCS:</b> J0897, J1740, J3110, J3111, J3489

\*Codes subject to change

Osteoporosis Medications		
Description	Prescription	(continued)



## (OSW) Osteoporosis Screening in Older Women *(continued)*

<b>Bisphosphonates</b>	• Alendronate • Ibandronate	• Alendronate-cholecalciferol • Risedronate • Zoledronic acid
<b>Other agents</b>	• Abaloparatide • Teriparatide	• Romosozumab • Denosumab • Raloxifene

### Action

- ✓ Make sure that all women ages 65-75 receive a routine osteoporosis screening with a BMD test

### Tips and Best Practices

- ✓ Support BMD scheduling with accessible in-network provider during patient visit and have referrals easily accessible or standing orders
- ✓ BMD tests are the most effective method for determining bone health, identifying osteoporosis, determining risk for fractures, and assessing response to osteoporosis treatment
- ✓ Review osteoporosis risk factors and long-term effects to reinforce importance of routine screening
- ✓ Provider tool kit and additional resource trainings are available through the Bone Health & Osteoporosis Foundation (BHOF) website at [www.bonehealthandosteoporosis.org/](http://www.bonehealthandosteoporosis.org/)

### Required Exclusions

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above or medical record documentation:

- ✓ Members who use hospice services any time during the measurement year

*(continued)*



## (OSW) Osteoporosis Screening in Older Women (continued)

- ✓ Members who die any time during the measurement year
- ✓ Members who received palliative care during the measurement year
- ✓ Members who had a dispensed prescription to treat osteoporosis any time between three years prior to the measurement year through December 31 of the year prior to the measurement year
- ✓ Members who had a claim/encounter for osteoporosis therapy any time in the member's history

.....



## (PPC) Prenatal and Postpartum Care

### Product Lines:

- ✓ Meridian (Medicaid)
- ✓ Ambetter from Meridian (Marketplace)

### Description

Measure evaluates percentage of deliveries of live births on or between October 8 of the year prior to the measurement year and October 7 of the measurement year. For these persons, the measure assesses the following facets of prenatal and postpartum care:

- ✓ **Timeliness of Prenatal Care:** percentage of deliveries that received a prenatal care visit in the first trimester, on or before the enrollment start date, or within 42 days of enrollment in the organization
- ✓ **Postpartum Care:** percentage of deliveries that had a postpartum visit on or between 7 and 84 days after delivery

Description	Codes*
-------------	--------



## (PPC) Prenatal and Postpartum Care

(continued)

Prenatal Visits <b>(must also include pregnancy related diagnosis code)</b>	<p><b>CPT:</b> 98000-98016, 98966-98968, 98970-98972, 98980, 98981, 99202-99205, 99211-99215, 99242-99245, 99421-99423, 99457, 99458, 99483, 99441-99443</p> <p><b>HCPCS:</b> G0071, G0463, G2010, G2012, G2250, G2251, G2252, T1015</p> <p><b>AND</b></p> <p><b>Pregnancy Diagnosis Code:</b> ICD-10: Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82, Z34.83, Z34.91, Z34.92, Z34.93</p> <p>*list does not include all codes for pregnancy, due to amount</p>
Prenatal Bundled Services	<p><b>CPT:</b> 59400, 59425, 52426, 59510, 59610, 59618</p> <p><b>HCPCS:</b> H1005</p>
Stand Alone Prenatal Visit (no dx codes needed)	<p><b>CPT-CAT-II:</b> 0500F, 0501F, 0502F</p> <p><b>CPT:</b> 99500</p> <p><b>HCPCS:</b> H100, H1001, H1002, H1003, H1004</p>
Postpartum Visits	<p><b>CPT:</b> 57170, 58300, 59430, 99501</p> <p><b>CPT-CAT-II:</b> 0503F</p> <p><b>HCPCS:</b> G0101</p> <p><b>ICD-10:</b> Z01.411, Z01.419, Z01.42, Z30.430, Z39.1, Z39.2</p>
Cervical Cytology	<p><b>CPT:</b> 88141-88143, 88147, 88148, 88150, 88152, 88153, 88164-88167, 88174, 88175</p> <p><b>HCPCS:</b> G0123, G0124, G0141, G0143-G0145, G0147, G0148, P3000, P3001</p>
Postpartum Bundled Services	<p><b>CPT:</b> 59400, 59410, 59510, 59515, 59610, 59614, 59618, 59622</p>
Hospice Care	<p><b>HCPCS:</b> G9473-G9479, Q5003-Q5008, Q5010, S9126, T2042-T2046, G0182</p> <p><b>CPT:</b> 99377, 99378</p>
Non-Live Births	<p><b>ICD-10:</b> O00.00, O00.01, O00.101, O00.102, O00.109, O00.111, O00.112, O00.119, O00.201, O00.202, O00.209</p>

\*Codes subject to change

(continued)



# (PPC) Prenatal and Postpartum Care

*(continued)*

## Action

- ✓ Ensure patients that have recently given birth are seen for postpartum care within 1-12 weeks of delivery with a qualified provider. Postpartum care should include one of the following, at a minimum:
  - Notation of “postpartum” care
  - Evaluation of weight, blood pressure, breasts, and abdomen
  - Perineal or cesarean incision/wound check
  - Screening for depression, anxiety, tobacco use, substance use, or preexisting mental health disorders
  - Glucose screening for women with gestational diabetes
  - Documentation of topics such as infant care or breastfeeding, resumption of intercourse, birth spacing or family planning, sleep/fatigue, or resumption of physical activity and healthy weight
- ✓ Ensure pregnant patients are seen for prenatal care in their first trimester with a qualified provider (OB/GYN, Midwife, PCP, Family Medicine, Family Practice, Nurse Practitioner or Physician Assistant). Prenatal care should include one of the following, at a minimum:
  - Diagnosis of pregnancy or reference to pregnancy
  - Documentation of LMP, EDD, or Gestational Age
  - Documentation of gravidity and parity
  - Positive pregnancy test
  - Basic OB exam that includes one of the following: auscultation for fetal heart tones, pelvic exam with OB observations or measurement of fundal height

*(continued)*



# (PPC) Prenatal and Postpartum Care

(continued)

## Tips and Best Practices

- ✓ Educate our mutual Medicaid members that six visits with a Doula are covered through our partner Mae Health. These can be anytime during their pregnancy through 1 year postpartum. Doulas are trained birth workers that give emotional and physical support to pregnant and postpartum individuals and their families. Having a doula during pregnancy has been shown to improve the health of birthing parents and their babies. Visit <https://meetmae.com/> to learn more!
- ✓ Visits do not need to be with an OB if a pregnancy diagnosis is included
- ✓ If member enrolls on our plan after first trimester, a prenatal visit should be completed within 42 days
- ✓ Telehealth visits, telephone visits, e-visits and virtual check-ins are acceptable for both prenatal and postpartum care
- ✓ Schedule postpartum visits prior to discharge after delivery
- ✓ Prenatal visits may occur inpatient as long as they are not the same admission as the delivery
- ✓ Make the most of the visit by providing other services that may be necessary during OB visit, such as chlamydia screening or cervical cancer screening

## Required Exclusions

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above or medical record documentation:

- ✓ Members who use hospice services any time during the measurement year
- ✓ Members who die any time during the measurement year
- ✓ Fetal demise/no live birth in the allowable time frame

---



# (PRS-E) Prenatal Immunization Status

## Product Lines:

- ✓ Meridian (Medicaid)

## Description

Measure evaluates the percentage of deliveries in the measurement year in which members had received influenza and tetanus, diphtheria toxoids and acellular pertussis (Tdap) vaccinations.

Description	Codes*
Adult Influenza Immunization	<b>CVX:</b> 88, 135, 140, 141, 144, 150, 153, 155, 158, 166, 168, 171, 185, 186, 197, 205, 320 <b>CPT:</b> 90653, 90656, 90658, 90661, 90662, 90673, 90674, 90682, 90686, 90688, 90689, 90694, 90756
Anaphylaxis Due To Influenza Vaccine	<b>SNOMED:</b> 471361000124100
Tdap Vaccine	<b>CVX:</b> 115 <b>CPT:</b> 90715
Anaphylaxis Due to Diphtheria, Tetanus or Pertussis Vaccine	<b>SNOMED:</b> 428281000124107, 428291000124105
Encephalitis Due to Diphtheria, Tetanus or Pertussis Vaccine	<b>SNOMED:</b> 192710009, 192711008, 192712001
Gestational Age Assessment <37 Weeks	<b>SNOMED:</b> 412726003 <b>ICD10CM:</b> Z3A.01, Z3A.08-Z3A.36
Hospice Care	<b>HCPCS:</b> G9473-G9479, Q5003-Q5008, Q5010, S9126, T2042-T2046, G0182 <b>CPT:</b> 99377, 99378

\*Codes subject to change

(continued)



# (PRS-E) Prenatal Immunization Status

(continued)

## Action

Perform the following immunizations in the listed timeframe:

- ✓ Influenza
  - Between July 1st of the year prior to the measurement year and the delivery date
- ✓ Tdap
  - During the pregnancy, including on the delivery date

## Tips and Best Practices

- ✓ If a member had anaphylaxis due to either of the required vaccines or encephalitis due to Tdap, submit codes in the table above to count towards the measure
- ✓ Enter administered immunizations in the Michigan Care Improvement Registry (MCIR). Information regarding MCIR can be found here: [\*\*https://mcir.org/training-resources/healthcare-provider-mcir-resources/\*\*](https://mcir.org/training-resources/healthcare-provider-mcir-resources/)
- ✓ Offer to administer immunizations at each prenatal visit until completed
- ✓ Offer immunizations on-site or support scheduling at accessible in-network facility during patient visit and have referrals easily accessible or standing orders
- ✓ Refer to the following website for CDC vaccine recommendations and clinical guidelines: [\*\*https://www.cdc.gov/vaccines-pregnancy/recommended-vaccines/index.html\*\*](https://www.cdc.gov/vaccines-pregnancy/recommended-vaccines/index.html)

## Required Exclusions

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above or medical record documentation:

- ✓ Deliveries that occurred at less than 37 weeks gestation



## (PRS-E) Prenatal Immunization Status (continued)

- ✓ Members who use hospice services any time during the measurement year
- ✓ Members who die any time during the measurement year

.....



## (PND-E) Prenatal Depression Screening and Follow-Up

### **Product Lines:**

- ✓ Meridian (Medicaid)

### **Description**

Measure evaluates the percentage of deliveries in the measurement year in which members were screened for clinical depression while pregnant and, if screened positive, received follow-up care.

### **Two rates are reported:**

1. Depression Screening: percentage of deliveries in which members were screened for clinical depression during pregnancy using a standardized instrument
2. Follow-Up on Positive Screen: percentage of deliveries in which members received follow-up care within 30 days of a positive depression screen finding

*(continued)*



## (PND-E) Prenatal Depression Screening and Follow-Up *(continued)*

Description	Codes*
<p><b>Follow-Up Visit (must have diagnosis of depression or other behavioral health condition)</b></p>	<p><b>Follow-Up Visit</b> <b>CPT:</b> 98000-98016, 98960-98962, 98966-98968, 98970-98972, 98980, 98981, 99078, 99202-99205, 99211-99215, 99242-99245, 99341-99345, 99347-99349, 99350, 99381-99387, 99391-99397, 99401-99404, 99411, 99412, 99421-99423, 99441-99443, 99457, 99458, 99483 <b>HCPCS:</b> G0071, G0463, G2010, G2012, G2250, G2251, G2252, T1015</p> <p><b>AND</b></p> <p><b>Depression or other BH Condition</b> <b>ICD10CM:</b> F01.51, F06.4, F11.188, F12.180, F12.980, F13.180, F14.180, F16.180, F20.0, F25.9, F31.3</p> <p>*list does not include all codes for BH conditions, due to amount</p>

*(continued)*



## (PND-E) Prenatal Depression Screening and Follow-Up *(continued)*

<p>Depression Case Management Encounter (<b>must</b> document assessment for symptoms of depression or dx of depression or other BH condition)</p>	<p><b>Depression Case Management Encounter</b>  <b>CPT:</b> 99366, 99492, 99493, 99494  <b>HCPCS:</b> G0512, T1016, T1017, T2022, T2023</p> <p><b>AND</b></p> <p><b>Symptoms of Depression</b>  <b>SNOMED:</b> 394924000, 788976000</p> <p><b>OR</b></p> <p><b>Depression or other BH Condition</b>  <b>ICD10CM:</b> F01.51, F06.4, F11.188, F12.180, F12.980, F13.180, F14.180, F16.180, F20.0, F25.9, F31.3  *list does not include all codes for BH conditions, due to amount</p>
<p><b>Behavioral Health Encounter</b></p>	<p><b>CPT:</b> 90791, 90792, 90832-90834, 90836-90839, 90845-90847, 90849, 90853, 90865, 90867-90870, 90875, 90876, 90880, 90887, 99484, 99492, 99493  <b>HCPCS:</b> G0155, G0176, G0177, G0409, G0410, G0411, G0511, G0512, H002, H004  *list does not include all codes for BH encounters, due to amount</p>
<p><b>Exercise Counseling</b></p>	<p><b>ICD10CM:</b> Z71.82</p>
<p><b>Gestational Age Assessment &lt;37 Weeks</b></p>	<p><b>SNOMED:</b> 412726003  <b>ICD10CM:</b> Z3A.01, Z3A.08-Z3A.36</p>
<p><b>Hospice Care</b></p>	<p><b>HCPCS:</b> G9473-G9479, Q5003-Q5008, Q5010, S9126, T2042-T2046, G0182  <b>CPT:</b> 99377, 99378</p>

*(continued)*



## (PND-E) Prenatal Depression Screening and Follow-Up (continued)

Depression Screening Instruments		
Instruments for Adolescents (≤17 years)	Total Score LOINC Codes	Positive Finding
Patient Health Questionnaire (PHQ-9)®	44261-6	Total score ≥ 10
Patient Health Questionnaire Modified for Teens (PHQ-9M)®	89204-2	Total score ≥ 10
Patient Health Questionnaire-2 (PHQ-2)®	55758-7	Total score ≥ 3
Beck Depression Inventory - Fast Screen (BDI-FS)®	89208-3	Total score ≥ 8
Center for Epidemiologic Studies Depression Scale - Revised (CESD-R)	89205-9	Total score ≥ 17
Edinburgh Postnatal Depression Scale (EPDS)	99046-5	Total score ≥ 10
PROMIS Depression	71965-8	Total score (T Score) ≥ 60

Depression Screening Instruments		
Instruments for Adolescents (18+years)	Total Score LOINC Codes	Positive Finding
Patient Health Questionnaire (PHQ-9)®	44261-6	Total score ≥ 10
Patient Health Questionnaire-2 (PHQ-2)®	55758-7	Total score ≥ 3
Beck Depression Inventory - Fast Screen (BDI-FS)®	89208-3	Total score ≥ 8

(continued)



## (PND-E) Prenatal Depression Screening and Follow-Up (continued)

Beck Depression Inventory (BDI-II) <sup>®</sup>	89209-1	Total score ≥ 20
Center for Epidemiologic Studies Depression Scale - Revised (CESD-R)	89205-9	Total score ≥ 17
Duke Anxiety-Depression Scale (DUKE-AD) <sup>®</sup>	90853-3	Total score ≥ 30
Edinburgh Postnatal Depression Scale (EPDS)	99046-5	Total score ≥ 10
My Mood Monitor (M-3) <sup>®</sup>	71777-7	Total score ≥ 5
PROMIS Depression	71965-8	Total score (T Score) ≥ 60
PROMIS Emotional Distress - Depression - Short Form	77861-3	Total score ≥ 60
Clinically Useful Depression Outcome Scale (CUDOS)	90221-3	Total score ≥ 31

\*Codes subject to change

### Action

- ✓ Screen all pregnant members for clinical depression using one of the standardized depression screening assessments in the table above
- ✓ For any positive screenings, follow-up within 30 days with one of the following:
  - Outpatient, telehealth, telephone, e-visit, or virtual-check in with a diagnosis of depression or other behavioral health condition
  - Depression case management encounter that documents assessment for symptoms of depression or a diagnosis of depression or other behavioral health condition
  - A behavioral health encounter, including assessment, therapy, collaborative care or medication management
  - Prescribe antidepressant medication
  - Exercise counseling
  - Documentation of additional depression screening on a



## (PND-E) Prenatal Depression Screening and Follow-Up *(continued)*

full-length instrument indicating either no depression or no symptoms that require follow-up on the same day as a positive screen on a brief screening instrument

### Tips and Best Practices

- ✓ If using a brief screening instrument (such as PHQ-2) results in a positive screen, immediately follow up with a full-length instrument (such as PHQ-9). If there are negative findings, this will be considered the follow-up
- ✓ Perform any of the specified prenatal depression screening tools at the first prenatal visit as part of your standard initial prenatal exam

### Required Exclusions

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above or medical record documentation:

- ✓ Deliveries that occurred at less than 37 weeks gestation
- ✓ Members who use hospice services any time during the measurement year
- ✓ Members who die any time during the measurement year

---



## (PDS-E) Postpartum Depression Screening and Follow-Up

### Product Lines:

- ✓ Meridian (Medicaid)

### Description

Measure evaluates the percentage of deliveries in the measurement year in which members were screened for clinical depression during the postpartum period, and, if screened positive, received follow-up care.



## (PDS-E) Postpartum Depression Screening and Follow-Up *(continued)*

### Two rates are reported:

1. Depression Screening: percentage of deliveries in which members were screened for clinical depression using a standardized instrument during the postpartum period
2. Follow-Up on Positive Screen: percentage of deliveries in which members received follow-up care within 30 days of a positive depression screen finding

DESCRIPTION	CODES*
<b>Follow-Up Visit (must have diagnosis of depression or other behavioral health condition)</b>	<b>Follow-Up Visit</b> <b>CPT:</b> 98000-98016, 98960-98962, 98966-98968, 98970-98972, 98980, 98981, 99078, 99202-99205, 99211-99215, 99242-99245, 99341-99345, 99347-99349, 99350, 99381-99387, 99391-99397, 99401-99404, 99411, 99412, 99421-99423, 99441-99443, 99457, 99458, 99483 <b>HCPCS:</b> G0071, G0463, G2010, G2012, G2250, G2251, G2252, T1015  <b>AND</b>  <b>Depression or other BH Condition</b> <b>ICD10CM:</b> F01.51, F06.4, F11.188, F12.180, F12.980, F13.180, F14.180, F16.180, F20.0, F25.9, F31.3 *list does not include all codes for BH conditions, due to amount



## (PDS-E) Postpartum Depression Screening and Follow-Up *(continued)*

DESCRIPTION	CODES*
Depression Case Management Encounter <b>(must document assessment for symptoms of depression or dx of depression or other BH condition)</b>	<p><b>Depression Case Management Encounter</b></p> <p><b>CPT:</b> 99366, 99492, 99493, 99494</p> <p><b>HCPCS:</b> G0512, T1016, T1017, T2022, T2023</p> <p><b>AND</b></p> <p><b>Symptoms of Depression</b></p> <p><b>SNOMED:</b> 394924000, 788976000</p> <p><b>OR</b></p> <p><b>Depression or other BH Condition</b></p> <p><b>ICD10CM:</b> F01.51, F06.4, F11.188, F12.180, F12.980, F13.180, F14.180, F16.180, F20.0, F25.9, F31.3</p> <p>*list does not include all codes for BH conditions, due to amount</p>
Behavioral Health Encounter	<p><b>CPT:</b> 90791, 90792, 90832-90834, 90836-90839, 90845-90847, 90849, 90853, 90865, 90867-90870, 90875, 90876, 90880, 90887, 99484, 99492, 99493</p> <p><b>HCPCS:</b> G0155, G0176, G0177, G0409, G0410, G0411, G0511, G0512, H002, H004</p> <p>*list does not include all codes for BH encounters, due to amount</p>
Exercise Counseling	<b>ICD10CM:</b> Z71.82
Hospice Care	<p><b>HCPCS:</b> G9473-G9479, Q5003-Q5008, Q5010, S9126, T2042-T2046, G0182</p> <p><b>CPT:</b> 99377, 99378</p>

\*Codes subject to change

*(continued)*



# (PDS-E) Postpartum Depression Screening and Follow-Up (continued)

Depression Screening Instruments		
Instruments for Adolescents (≤17 years)	Total Score LOINC Codes	Positive Finding
Patient Health Questionnaire (PHQ-9)®	44261-6	Total score ≥ 10
Patient Health Questionnaire Modified for Teens (PHQ-9M)®	89204-2	Total score ≥ 10
Patient Health Questionnaire-2 (PHQ-2)®	55758-7	Total score ≥ 3
Beck Depression Inventory - Fast Screen (BDI-FS)®	89208-3	Total score ≥ 8
Center for Epidemiologic Studies Depression Scale - Revised (CESD-R)	89205-9	Total score ≥ 17
Edinburgh Postnatal Depression Scale (EPDS)	99046-5	Total score ≥ 10
PROMIS Depression	71965-8	Total score (T Score) ≥ 60

Depression Screening Instruments		
Instruments for Adolescents (18+years)	Total Score LOINC Codes	Positive Finding
Patient Health Questionnaire (PHQ-9)®	44261-6	Total score ≥ 10
Patient Health Questionnaire-2 (PHQ-2)®	55758-7	Total score ≥ 3
Beck Depression Inventory - Fast Screen (BDI-FS)®	89208-3	Total score ≥ 8
Beck Depression Inventory (BDI-II)®	89209-1	Total score ≥ 20

(continued)



# (PDS-E) Postpartum Depression Screening and Follow-Up (continued)

Center for Epidemiologic Studies Depression Scale - Revised (CESD-R)	89205-9	Total score $\geq$ 17
Duke Anxiety-Depression Scale (DUKE-AD) <sup>®</sup>	90853-3	Total score $\geq$ 30
Edinburgh Postnatal Depression Scale (EPDS)	71354-5	Total score $\geq$ 10
My Mood Monitor (M-3) <sup>®</sup>	71777-7	Total score $\geq$ 5
PROMIS Depression	71965-8	Total score (T Score) $\geq$ 60
PROMIS Emotional Distress - Depression - Short Form	77861-3	Total score $\geq$ 60
Clinically Useful Depression Outcome Scale (CUDOS)	90221-3	Total score $\geq$ 31

## Action

- ✓ Screen all members within 7-84 days after delivery for clinical depression using one of the standardized depression screening assessments in the table above
- ✓ For any positive screenings, follow-up within 30 days with one of the following:
  - Outpatient, telehealth, telephone, e-visit, or virtual-check in with a diagnosis of depression or other behavioral health condition
  - Depression case management encounter that documents assessment for symptoms of depression or a diagnosis of depression or other behavioral health condition
  - A behavioral health encounter, including assessment, therapy, collaborative care or medication management
  - Prescribe antidepressant medication
  - Exercise counseling
  - Documentation of additional depression screening on a full-length instrument indicating either no depression or no symptoms that require follow-up on the same day as a positive screen on a brief screening instrument



# (PDS-E) Postpartum Depression Screening and Follow-Up *(continued)*

## Tips and Best Practices

- ✓ If using a brief screening instrument (such as PHQ-2) results in a positive screen, immediately follow up with a full-length instrument (such as PHQ-9). If there are negative findings, this will be considered the follow-up
- ✓ Perform any of the specified postpartum depression screening tools as part of your standard postpartum exam

## Required Exclusions

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above or medical record documentation:

- ✓ Members who use hospice services any time during the measurement year
- ✓ Members who die any time during the measurement year

.....

# Pediatric Health

---



## (ADD-E) Follow-Up Care for Children Prescribed ADHD Medication

### Product Lines:

- ✓ Meridian (Medicaid)

### Description

Measure evaluates percentage of children 6-12 years of age newly prescribed attention-deficit/hyperactivity disorder (ADHD) medication who had at least three follow up care visits with a practitioner with prescribing authority:

- ✓ At least one follow-up visit within 30 days of prescription dispense date
- ✓ At least two follow-up visits 1-9 months after prescription dispense date

Description	Codes*
BH Outpatient Visit	<b>CPT:</b> 98960-98962, 99078, 99202-99205, 99211-99215, 99242-99245, 99341-99345, 99347-99350, 99381-99387, 99391-99397, 99401-99404, 99411, 99412, 99483, 99492, 99493, 99494, 99510 <b>HCPCS:</b> G0155, G0176, G0177, G0409, G0468, G0512, H0002, H0004, H0031, H0034, H0036, H0037, H0039, H0040, H2000, H2010, H2011, H2013, H2014-H2020, T1015

(continued)



## (ADD-E) Follow-Up Care for Children Prescribed ADHD Medication (continued)

Description	Codes*
<b>Visit with unspecified setting (must also include POS code)</b>	<b>CPT:</b> 90791, 90792, 90832-90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99252-99255 <b>AND</b> <b>POS:</b> 02, 03, 05, 07, 09, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 22, 33, 49, 50, 52, 53, 71, 72
<b>Health and Behavior Assessment or Intervention</b>	<b>CPT:</b> 96156, 96158, 96159, 96164, 96168, 96170, 96171
<b>Intensive Outpatient Encounter or Partial Hospitalization</b>	<b>HCPCS:</b> G0410, G0411, H0035, H2001, H2012, S0201, S9480, S9484, S9485
<b>Telephone Visits</b>	<b>CPT:</b> 98966, 98967, 98968, 99441, 99442, 99443
<b>E-visit or virtual check-in</b>	<b>CPT:</b> 98970, 98971, 98972, 98980, 98981, 99421, 99422, 99423, 99457, 99458 <b>HCPCS:</b> G0071, G2010, G2012, G2250, G2251, G2252
<b>Narcolepsy</b>	<b>ICD10CM:</b> G47.411, G47.419, G47.421, G47.429
<b>Hospice Care</b>	<b>HCPCS:</b> G9473-G9479, Q5003-Q5008, Q5010, S9126, T2042-T2046, G0182 <b>CPT:</b> 99377, 99378

\*Codes subject to change

### Action

- ✓ Schedule a follow-up appointment within one month for patients newly prescribed ADHD medication. Schedule a follow-up visit while your patient is still in the office
- ✓ Schedule two or more visits in the nine months after the initial follow-up visit to continue to monitor your patient's progress

(continued)



## (ADD-E) Follow-Up Care for Children Prescribed ADHD Medication (continued)

### Tips and Best Practices

- ✓ Offer flexible appointment availability including evening, weekends, and telehealth options

### Required Exclusions

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above or medical record documentation:

- ✓ Members who use hospice services any time during the measurement year
- ✓ Members who die any time during the measurement year
- ✓ Members with a diagnosis of narcolepsy any time during the member's history

---



## (APM-E) Metabolic Monitoring for Children and Adolescents on Antipsychotics

### Product Lines:

- ✓ Meridian (Medicaid)

### Description

Measure evaluates the percentage of children and adolescents 1-17 years of age who had two or more antipsychotic prescriptions and had metabolic testing. Three rates reported:

- 1 Percentage of children and adolescents on antipsychotics who received blood glucose testing
- 2 Percentage of children and adolescents on antipsychotics who received cholesterol testing
- 3 Percentage of children and adolescents on antipsychotics who received blood glucose and cholesterol testing

(continued)



## (APM-E) Metabolic Monitoring for Children and Adolescents on Antipsychotics (continued)

Description (Need either A1c or Glucose and LCL-C or Cholesterol)	Codes*
HbA1c Lab Tests	<b>CPT:</b> 83036, 83037 <b>CPT-CAT-II:</b> 3044F, 3046F, 3051F, 3052F
Glucose Lab Tests	<b>CPT:</b> 80047, 80048, 80050, 80053, 80069, 82947, 82950, 82951
LDL-C Lab Tests	<b>CPT:</b> 80061, 83700, 83701, 83704, 83721 <b>CPT-CAT-II:</b> 3048F, 3049F, 3050F
Cholesterol Lab Tests	<b>CPT:</b> 82465, 83718, 83722, 84478
Hospice Care	<b>HCPCS:</b> G9473-G9479, Q5003-Q5008, Q5010, S9126, T2042-T2046, G0182 <b>CPT:</b> 99377, 99378

\*Codes subject to change

### Action

- ✓ Ensure patients who had two or more antipsychotic prescriptions have the following testing completed in the measurement year have:
  - A blood glucose test **or** HbA1C
  - A cholesterol **or** LDL-C test
- ✓ These tests can be on the same or different dates of service

### Tips and Best Practices

- ✓ Provide members/caregivers with lab orders for HbA1c or glucose and cholesterol or LDL-C to be completed yearly
- ✓ Educate the member and caregiver about the risks associated with taking antipsychotic medications and the importance of regular follow up care

(continued)



## (APM-E) Metabolic Monitoring for Children and Adolescents on Antipsychotics (continued)

- ✓ Offer lab testing on-site or support lab scheduling at accessible in-network facility during patient visits and have referrals easily accessible or standing orders

### Required Exclusions

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above or medical record documentation:

- ✓ Members who use hospice services any time during the measurement year
- ✓ Members who die any time during the measurement year

---



## (CIS-E) Childhood Immunization Status

### Product Lines:

- ✓ Meridian (Medicaid)
- ✓ Ambetter from Meridian (Marketplace)

### Description

Measure evaluates the percentage of children who turned two during the measurement year and completed the following immunizations on or before their second birthday:

- ✓ 4 DTAP (diphtheria, tetanus, and acellular pertussis) **or** anaphylaxis **or** encephalitis due to DTAP
- ✓ 3 IPV (polio) **or** anaphylaxis due to IPV
- ✓ 1 MMR (measles, mumps and rubella) **or** history of measles illness, mumps illness, and rubella illness **or** anaphylaxis due to MMR

(continued)



## (CIS-E) Childhood Immunization Status

(continued)

- ✓ 3 HIB (haemophilus influenza type B) **or** anaphylaxis due to HiB
- ✓ 3 HepB (hepatitis B) **or** anaphylaxis due to HepB **or** history of HepB illness
- ✓ 1 VZV (chicken pox) **or** history of chicken pox illness **or** anaphylaxis due to VZV
- ✓ 4 PCV (pneumococcal conjugate) **or** anaphylaxis due to PCV
- ✓ 1 HepA (hepatitis A) **or** history of HepA illness **or** anaphylaxis due to HepA
- ✓ 2 or 3 RV (rotavirus) **or** anaphylaxis due to RV
- ✓ 2 FLU (influenza) **or** anaphylaxis due to FLU vaccine

Description	Codes*
DTAP Immunization	<b>CPT:</b> 90697, 90698, 90700, 90723 <b>CVX:</b> 20, 50, 106, 107, 110, 120, 146, 198
Anaphylaxis Due to DTAP	<b>SNOMED:</b> 428281000124107, 428291000124105
Encephalitis Due to DTAP	<b>SNOMED:</b> 192710009, 192711008, 192712001
HIB Immunization	<b>CPT:</b> 90644, 90647, 90648, 90697, 90698, 90748 <b>CVX:</b> 17, 46-51, 120, 146, 148, 198
Anaphylaxis due to HIB	<b>SNOMED:</b> 433621000124101
Hep B Immunization	<b>CPT:</b> 90697, 90723, 90740, 90744, 90747, 90748 <b>CVX:</b> 08, 44, 45, 51, 110, 146, 198 <b>HCPCS:</b> G0010
Newborn Hep B Vaccine Administered	<b>ICD10PCS:</b> 3E0234Z
Hep B Illness	<b>ICD-10:</b> B16.0, B16.1, B16.2, B16.9, B17.0, B18.0, B18.1, B19.10, B19.11
Anaphylaxis due to HepB	<b>SNOMED:</b> 428321000124101
IPV Immunization	<b>CPT:</b> 90697, 90698, 90713, 90723 <b>CVX:</b> 10, 89, 110, 120, 146



## (CIS-E) Childhood Immunization Status

(continued)

Description	Codes*
Anaphylaxis due to IPV	<b>SNOMED:</b> 471321000124106
MMR Immunization	<b>CPT:</b> 90707, 90710 <b>CVX:</b> 03, 94
Anaphylaxis due to MMR	<b>SNOMED:</b> 471331000124109
Measles, Mumps, and Rubella Illness	<b>Measles, ICD-10:</b> B05.0-B05.4, B05.81, B05.89, B05.9 <b>Mumps, ICD-10:</b> B26.0, B26.1, B26.2, B26.3, B26.81, B26.82, B26.83, B26.84, B26.85, B26.89, B26.9 <b>Rubella, ICD-10:</b> B06.00, B06.01, B06.02, B06.09, B06.81, B06.82, B06.89, B06.9
PCV Immunization	<b>CPT:</b> 90670, 90671, 90677 <b>CVX:</b> 109, 133, 152, 215, 216 <b>HCPCS:</b> G0009
Anaphylaxis due to PCV	<b>SNOMED:</b> 471141000124102
VZV Immunization	<b>CPT:</b> 90710, 90716 <b>CVX:</b> 21, 94
VZV (Chicken Pox) Illness	<b>ICD-10:</b> B01.0, B01.11, B01.12, B01.2, B01.81, B01.89, B01.9, B02.0, B02.1, B02.21, B02.22, B02.23, B02.24, B02.29, B02.30, B02.31, B02.32, B02.33, B02.34, B02.39, B02.7, B02.8, B02.9
Anaphylaxis due to VZV	<b>SNOMED:</b> 471341000124104
Hep A Immunization	<b>CPT:</b> 90633 <b>CVX:</b> 31, 83, 85
HepA Illness	<b>ICD-10:</b> B15.0, B15.9
Anaphylaxis due to HepA	<b>SNOMED:</b> 471311000124103
Influenza Immunization	<b>CPT:</b> 90655-90658, 90661, 90685-90689, 90674, 90756 <b>CVX:</b> 88, 140, 141, 150, 153, 155, 158, 161, 171, 186, 320 <b>HCPCS:</b> G0008

(continued)



## (CIS-E) Childhood Immunization Status

(continued)

Description	Codes*
Anaphylaxis due to Influenza	<b>SNOMED:</b> 471361000124100
Rotavirus 2-Dose Immunization	<b>CPT:</b> 90681 <b>CVX:</b> 119
Rotavirus 3-Dose Immunization	<b>CPT:</b> 90680 <b>CVX:</b> 116, 122
Anaphylaxis due to Rotavirus	<b>SNOMED:</b> 428331000124103
Organ and Bone Marrow Transplants	<b>CPT:</b> 32850-32856, 33927-33930, 33933, 33935, 33940, 33944, 33945, 44132, 44133, 44135-44137, 44715, 44720, 44721, 47133, 47135, 47136, 47140-47147, 48160, 48550-48554, 48556, 50360, 50365, 50380
Hospice Care	<b>HCPCS:</b> G9473-G9479, Q5003-Q5008, Q5010, S9126, T2042-T2046, G0182 <b>CPT:</b> 99377, 99378

\*Codes subject to change

### Action

Perform the following immunizations in listed timeframe:

- ✓ **From birth through second birthday**
  - Hepatitis B
- ✓ **From 42 days after birth through second birthday**
  - DTAP, IPV, HIB, PCV, Rotavirus
- ✓ **From 6 months through second birthday**
  - Influenza
- ✓ **From first birthday through second birthday**
  - MMR, VZV, Hepatitis A

(continued)



# (CIS-E) Childhood Immunization Status

(continued)

## Tips and Best Practices

- ✓ Enter administered immunizations in the Michigan Care Improvement Registry (MCIR). Information regarding MCIR can be found here: [\*\*https://mcir.org/training-resources/healthcare-provider-mcir-resources/\*\*](https://mcir.org/training-resources/healthcare-provider-mcir-resources/)
- ✓ If newborn Hep B vaccination is administered, ensure it is coded in the child's record
- ✓ Rotavirus can be administered in a 2 dose (RotaTeq or RV5) or 3 dose (Rotarix or RV1) series. 1 dose of RV5 and 2 doses of RV1 can combine to count towards compliance for the 3 dose series
- ✓ Parent/caregiver reported vaccines are acceptable if documented in chart with vaccine name and administration date
- ✓ Try to complete immunizations at scheduled well-exams to reduce the number of appointments
- ✓ If you are not currently offering immunizations at your office, check out the Vaccines for Children Program and how to get vaccines at no cost to your office! Contact your local health department or visit the Michigan VFC resource guide for details at [\*\*www.Michigan.gov/VFC\*\*](http://www.Michigan.gov/VFC)
- ✓ Educate patients and parents/guardians on the diseases that are prevented with immunizations
- ✓ Address concerns and fears that parents may have regarding vaccination
- ✓ Recommend the following reputable health resources for parent-friendly information and answers to common questions
  - Centers for Disease Control & Prevention (CDC) website at [\*\*https://www.cdc.gov/vaccines-children/index.html\*\*](https://www.cdc.gov/vaccines-children/index.html)
  - Michigan Department of Health and Human Services (MDHHS) website at [\*\*ivaccinate.org/answering-your-questions\*\*](http://ivaccinate.org/answering-your-questions)



## (CIS-E) Childhood Immunization Status

(continued)

### Required Exclusions

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above or medical record documentation:

- ✓ Members who had a contraindication to a childhood vaccine on or before their second birthday
- ✓ Members who had an organ or bone marrow transplant
- ✓ Members who use hospice services any time during the measurement year
- ✓ Members who die any time during the measurement year

---



## (IMA-E) Immunizations for Adolescents

### Product Lines

- ✓ Meridian (Medicaid)
- ✓ Ambetter from Meridian (Marketplace)

### Description

Measure evaluates the percentage of adolescents who turned 13 during the measurement year and completed the following immunizations on or before their 13th birthday:

- ✓ 1 Meningococcal **or** anaphylaxis due to Meningococcal
- ✓ 1 Tdap (tetanus, diphtheria toxoids and acellular pertussis) **or** anaphylaxis **or** encephalitis due to Tdap
- ✓ 2 or 3 HPV (human papillomavirus) **or** anaphylaxis due to HPV

(continued)



## (IMA-E) Immunizations for Adolescents

(continued)

Description	Codes*
Meningococcal Immunization	<b>CPT:</b> 90619, 90623, 90733, 90734 <b>CVX:</b> 32, 108, 114, 136, 147, 167, 203, 316, 328
Anaphylaxis due to Meningococcal Vaccine	<b>SNOMED:</b> 428301000124106
Tdap Immunization	<b>CPT:</b> 90715 <b>CVX:</b> 115
Anaphylaxis due to Tdap Vaccine	<b>SNOMED:</b> 428281000124107, 428291000124105
Encephalitis due to Tdap Vaccine	<b>SNOMED:</b> 192710009, 192711008, 192712001
HPV Immunization	<b>CPT:</b> 90649-90651 <b>CVX:</b> 62, 118, 137, 165
Anaphylaxis due to HPV Vaccine	<b>SNOMED:</b> 428241000124101
Hospice Care	<b>HCPCS:</b> G9473-G9479, Q5003-Q5008, Q5010, S9126, T2042-T2046, G0182 <b>CPT:</b> 99377, 99378

\*Codes subject to change

### Action

Perform the following immunizations in listed timeframe:

- ✓ From 9th through 13th birthday
  - HPV
- ✓ From 10th through 13th birthday
  - Tdap
  - Meningococcal

(continued)



## (IMA-E) Immunizations for Adolescents

(continued)

### Tips and Best Practices

- ✓ Enter administered immunizations in the Michigan Care Improvement Registry (MCIR). Information regarding MCIR can be found here: [\*\*https://mcir.org/training-resources/healthcare-provider-mcir-resources/\*\*](https://mcir.org/training-resources/healthcare-provider-mcir-resources/)
- ✓ If HPV vaccines are administered less than 146 days apart, 3 doses will be needed rather than 2
- ✓ Parent/caregiver reported vaccines are acceptable if documented in chart with vaccine name and administration date. Ensure your EMR has appropriate fields to house this data for all visit types.
- ✓ Try to complete immunizations at scheduled well-exams to reduce the number of appointments
- ✓ If you are not currently offering immunizations at your office, check out the Vaccines for Children Program and how to get vaccines at no cost to your office! Contact your local health department or visit the Michigan VFC resource guide for details at [\*\*www.Michigan.gov/VFC\*\*](http://www.Michigan.gov/VFC)
- ✓ Advise that the HPV vaccination is preventive. Although their child may not currently be sexually active, it is important to receive the HPV vaccination now to prevent their child from getting HPV in the future
- ✓ Recommend the following reputable health resources for parent-friendly information and answers to common questions:
  - Centers for Disease Control & Prevention (CDC) website at [\*\*https://www.cdc.gov/vaccines-children/index.html\*\*](https://www.cdc.gov/vaccines-children/index.html)
  - Michigan Department of Health and Human Services (MDHHS) website at [\*\*ivaccinate.org/answering-your-questions\*\*](http://ivaccinate.org/answering-your-questions)

(continued)



## (IMA-E) Immunizations for Adolescents

(continued)

### Required Exclusions

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above or medical record documentation:

- ✓ Members who use hospice services any time during the measurement year
- ✓ Members who die any time during the measurement year

.....



## (LSC-E) Lead Screening in Children

### Product Lines

- ✓ Meridian (Medicaid)

### Description

Measure evaluates the percentage of children who turned 2 during the measurement year who had one or more capillary or venous lead blood test for lead poisoning by their second birthday.

Description	Codes*
Lead Tests	<b>CPT:</b> 83655 <b>LOINC:</b> 10368-9, 10912-4, 14807-2, 17052-2, 25459-9, 27129-6, 32325-3, 5674-7, 77307-7
Hospice Care	<b>HCPCS:</b> G9473-G9479, Q5003-Q5008, Q5010, S9126, T2042-T2046, G0182 <b>CPT:</b> 99377, 99378

\*Codes subject to change



## (LSC-E) Lead Screening in Children (continued)

### Action

- ✓ Perform capillary or venous blood lead screening for all children on or before their second birthday

### Tips and Best Practices

- ✓ While HEDIS only requires one test prior to 24 months of age, providers should test at 12 and 24 months to be adherent to Michigan regulatory requirements and guidance
- ✓ Member-reported lead screenings are acceptable when documented in chart with test date and result. Ensure your EMR has appropriate fields to house this data for all visit types.
- ✓ Offer testing on-site or support scheduling at accessible in-network facility during patient visit and have referrals easily accessible or standing orders
- ✓ Enter lead screening results in the Michigan Care Improvement Registry (MCIR)
- ✓ Reach out to your Quality or Provider Network Management Representative to find out about Point of Care Lead Analyzers for your office
- ✓ For more information about blood lead testing, state guidance, and how to support patients with high blood lead levels, please visit **[www.michigan.gov/mileadsafe/professionals/healthcare-providers](http://www.michigan.gov/mileadsafe/professionals/healthcare-providers)**

### Required Exclusions

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above or medical record documentation:

- ✓ Members who use hospice services any time during the measurement year
- ✓ Members who die any time during the measurement year



## (OED) Oral Evaluation, Dental Services

### Product Lines:

- ✓ Meridian (Medicaid)
- ✓ Ambetter from Meridian (Marketplace)

### Description

Measure evaluates the percentage of members under 21 years of age who received a comprehensive or periodic oral evaluation with a dental provider during the measurement year.

Description	Codes*
Oral Evaluation	<b>CDT:</b> D0120, D0145, D0150
Hospice Care	<b>HCPCS:</b> G9473-G9479, Q5003-Q5008, Q5010, S9126, T2042-T2046, G0182 <b>CPT:</b> 99377, 99378

\*Codes subject to change

### Action

- ✓ Dental practitioner should perform dental visit at least once per year to prevent disease, reverse disease processes, prevent progression of caries, and reduce incidence of future lesions
- ✓ Dental practitioner is defined as:
  - Doctor of Dental Surgery (DDS)
  - Doctor of Dental Medicine (DMD)
  - Certified and Licensed Dental Hygienist

### Tips and Best Practices

- ✓ Encourage new patients to establish a dental home with your practice to ensure good routine oral healthcare and follow ups
- ✓ Send parents reminders every six months to schedule for periodic exams, prophylaxis (cleanings), and fluoride treatments

(continued)



## (OED) Oral Evaluation, Dental Services (continued)

- ✓ Provide local primary care practitioners with your practice's contact information for easy member referral
- ✓ Remind expectant mothers to make dental appointments for the baby either at the eruption of the first tooth or by the age of one

### Required Exclusions

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above or medical record documentation:

- ✓ Members who use hospice services any time during the measurement year
- ✓ Members who die any time during the measurement year

.....



## (TFC) Topical Fluoride for Children

### Product Lines:

- ✓ Meridian (Medicaid)

### Description

Measure evaluates the percentage of members 1-4 years of age who received at least two fluoride varnish applications during the measurement year.

Description	Codes*
Application of Fluoride Varnish	<b>CPT:</b> 99188 <b>CDT:</b> D1206

(continued)



## (TFC) Topical Fluoride for Children (continued)

### Hospice Care

**HCPCS:** G9473-G9479, Q5003-Q5008, Q5010,

S9126, T2042-T2046, G0182

**CPT:** 99377, 99378

\*Codes subject to change

### Action

- ✓ Pediatrician or dental practitioner should provide at least two fluoride varnish applications each year to prevent tooth decay

### Tips and Best Practices

- ✓ As soon as first tooth erupts, provide preventive fluoride varnish application at least once every six months
- ✓ Providers should recommend use of fluoridated toothpaste and amounts to be used based on age, provide educational materials, and ensure medical records include date that topical fluoride was applied and that it was coded appropriately
- ✓ Encourage parents to establish a dental home for their child to ensure good routine oral healthcare and follow ups

### Required Exclusions

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above or medical record documentation:

- ✓ Members who use hospice services any time during the measurement year
- ✓ Members who die any time during the measurement year



# (WCV) Child and Adolescent Well-Care Visits

## Product Lines

- ✓ Meridian (Medicaid)
- ✓ Ambetter from Meridian (Marketplace)

## Description

Measure evaluates the percentage of members 3-21 years of age who had at least one comprehensive well-care visit with a PCP or an OB/GYN practitioner during the measurement year.

Description	Codes*
Well-Care Visits	<b>CPT:</b> 99381-99385, 99391-99395, 99461 <b>HCPCS:</b> G0438, G0439, S0302, S0610, S0612, S0613 <b>ICD-10:</b> Z00.00, Z00.01, Z00.110, Z00.111, Z00.121, Z00.129, Z00.2, Z00.3, Z01.411, Z01.419, Z02.5, Z76.1, Z76.2, Z02.84
Hospice Care	<b>HCPCS:</b> G9473-G9479, Q5003-Q5008, Q5010, S9126, T2042-T2046, G0182 <b>CPT:</b> 99377, 99378

\*Codes subject to change

## Action

- ✓ Perform annual checkups to evaluate physical, developmental, behavioral, and emotional well-being.
- ✓ Conduct periodic Health Care Transition readiness/self-care skills assessments and provider anticipatory guidance and health education to assist members gain needed self-care skills

## Tips and Best Practices

- ✓ When a child is in your office for a sick visit or back-to-school/sports physical, also conduct a well-child visit if appropriate.

(continued)



## (WCV) Child and Adolescent Well-Care Visits *(continued)*

- ✓ Screening and counseling for obesity should occur at these visits by calculating a child's Body Mass Index (BMI) percentile for gender and age or plotting the value on a growth curve. Discuss physical activity and nutrition at every visit
- ✓ While a patient is in your office for a well-child visit, administer required vaccinations and testing

### Required Exclusions

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above or medical record documentation:

- ✓ Members who use hospice services any time during the measurement year
- ✓ Members who die any time during the measurement year

---



## (W30) Well Child Visits in the First 30 Months of Life

### Product Lines

- ✓ Meridian (Medicaid)
- ✓ Ambetter from Meridian (Marketplace)

### Description

Measure evaluates the percentage of members who turned 15 or 30 months old during the measurement year and who had:

*(continued)*



## (W30) Well Child Visits in the First 30 Months of Life (continued)

- ✓ **15 months:** 6 or more well-child visits between the ages of 0-15 months
- ✓ **30 months:** two or more well-child visits between the ages of 15-30 months

Description	Codes*
Well-Care Visits	<b>CPT:</b> 99381–99385, 99391–99395, 99461 <b>HCPCS:</b> G0438, G0439, S0302, S0610, S0612, S0613 <b>ICD-10:</b> Z00.00, Z00.01, Z00.110, Z00.111, Z00.121, Z00.129, Z00.2, Z00.3, Z01.411, Z01.419, Z02.5, Z76.1, Z76.2
Hospice Care	<b>HCPCS:</b> G9473–G9479, Q5003–Q5008, Q5010, S9126, T2042–T2046, G0182 <b>CPT:</b> 99377, 99378

\*Codes subject to change

### Action

- ✓ Complete appointments with all assigned patients according to the Bright Futures/American Academy of Pediatrics (AAP) Recommendations for Preventive Pediatric Health Care Periodicity Schedule

### Tips and Best Practices

- ✓ When a child is in your office for a sick visit, also conduct a well-child visit if appropriate.
- ✓ While a patient is in your office for a well-child visit, administer required vaccinations and testing
- ✓ Schedule next appointment at close of visit and offer flexible appointment availability including evening, weekends, telehealth, and family appointments



## (W30) Well Child Visits in the First 30 Months of Life *(continued)*

### Required Exclusions

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above or medical record documentation:

- ✓ Members who use hospice services any time during the measurement year
- ✓ Members who die any time during the measurement year

.....



## (WCC) Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents

### Product Lines

- ✓ Meridian (Medicaid)
- ✓ Ambetter from Meridian (Marketplace)

### Description

Measure evaluates the percentage of members 3-17 years of age who had an outpatient visit with a PCP or OB/GYN and who had evidence of the following:

- ✓ BMI Percentile documentation
- ✓ Counseling for Nutrition
- ✓ Counseling for Physical Activity

Description	Codes*
BMI Percentile	<b>ICD-10:</b> Z68.51, Z68.52, Z68.53, Z68.54, Z68.55, Z68.56

*(continued)*



## (WCC) Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents (continued)

<b>Nutrition Counseling</b>	<b>CPT:</b> 97802, 97803, 97804 <b>HCPCS:</b> G0270, G0271, G0447, S9449, S9452, S9470
<b>Physical Activity Counseling</b>	<b>HCPCS:</b> G0447, S9451 <b>ICD-10:</b> Z02.5, Z71.82
<b>Pregnancy Diagnosis</b>	<b>ICD-10:</b> Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82, Z34.83, Z34.91, Z34.92, Z34.93 *list does not include all codes for pregnancy, due to amount
<b>Hospice Care</b>	<b>HCPCS:</b> G9473-G9479, Q5003-Q5008, Q5010, S9126, T2042-T2046, G0182 <b>CPT:</b> 99377, 99378

\*Codes subject to change

### Action

- ✓ Complete a BMI percentile assessment at any visit for patients between 3 and 17 years of age. BMI percentile should be calculated or plotted on an appropriate age-growth chart
- ✓ Complete a nutrition and physical activity assessment followed by counseling or anticipatory guidance at any visit for patients 3 to 17 years of age

### Tips and Best Practices

- ✓ This measure couples well with well-child visits. Both measures will benefit from one visit. Please ensure correct coding when billing for both measures
- ✓ Utilize sick visits and sports physicals to also complete this measure. Include and document all three measure components during a sick visit for a compliant WCC record
- ✓ When counseling for nutrition, discuss appropriate food intake, healthy eating habits, issues including body image and eating disorders, etc.

(continued)



## **(WCC) Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents (continued)**

- ✓ When counseling for physical activity, discuss organized sports, activities, and document age-appropriate activity such as “rides bike for 30 minutes a day”
- ✓ Member-reported values for height, weight, and BMI percentile are acceptable when documented in chart

### **Required Exclusions**

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above or medical record documentation:

- ✓ Members who use hospice services any time during the measurement year
- ✓ Members who die any time during the measurement year
- ✓ Members who have a diagnosis of pregnancy during the measurement year

---

# General Health

---



## (CWP) Appropriate Testing for Pharyngitis

### Product Lines

- ✓ Meridian (Medicaid)
- ✓ Ambetter from Meridian (Marketplace)
- ✓ WellCare (Medicare)
- ✓ MI Coordinated Health (Dual Eligible Plan)

### Description

This measure demonstrates the percentage of episodes for members 3 years and older where the member was diagnosed with pharyngitis, dispensed an antibiotic, and received a group A streptococcus (strep) test for the episode.

Description	Codes*
Group A Strep Tests	<b>CPT:</b> 87070, 87071, 87081, 87430, 87650, 87651, 87652, 87880
Hospice Care	<b>HCPCS:</b> G9473-G9479, Q5003-Q5008, Q5010, S9126, T2042-T2046, G0182 <b>CPT:</b> 99377, 99378

\*Codes subject to change

### Action

- ✓ Perform a rapid strep test or throat culture to confirm diagnosis before prescribing antibiotics

### Tips and Best Practices

- ✓ Educate patients that an antibiotic is not necessary for viral infections, if rapid strep test and/or throat culture is negative
- ✓ Offer non-antibiotic symptom relief options that can provide comfort for viral infections

*(continued)*



## (CWP) Appropriate Testing for Pharyngitis

(continued)

- ✓ The Center for Disease Control and Prevention's Be Antibiotics Aware campaign offers many resources and trainings to help educate patients on appropriateness of antibiotic use on their website at [www.cdc.gov/antibiotic-use/index.html](http://www.cdc.gov/antibiotic-use/index.html)

### Required Exclusions

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above or medical record documentation:

- ✓ Member who use hospice services any time during the measurement year
- ✓ Member who die any time during the measurement year

---



## (FUA) Follow-Up After Emergency Department Visit for Substance Use

### Product Lines:

- ✓ WellCare (Medicare)
- ✓ MI Coordinated Health (Dual Eligible Plan)

### Description

Measure evaluates percentage of emergency department (ED) visits for members 13 years of age and older with a principal diagnosis of substance use disorder (SUD), or any diagnosis of drug overdose, and had a follow-up service.

(continued)



## (FUA) Follow-Up After Emergency Department Visit for Substance Use (continued)

Two rates are reported:

- ✓ ED visits for which the member received follow-up within 30 days of discharge
- ✓ ED visits for which the member received follow-up within 7 days of discharge

Description	Codes*
<b>Outpatient Visit</b> *must be with mental health provider or include appropriate dx codes	<b>CPT:</b> 90791, 90792, 90832-90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99252-99255  <b>Include Outpatient POS code from following:</b> POS: 03, 05, 07, 09, 11-20, 22, 33, 49, 50, 71, 72
<b>BH Outpatient Visit</b>	<b>CPT:</b> 98000-98007, 98960-98962, 99078, 99202-99205, 99211-99215, 99242-99245, 99341-99345, 99347-99350, 99381-99387, 99391-99397, 99401-99404, 99411, 99412, 99483, 99492-99494, 99510 <b>HCPCS:</b> G0155, G0176, G0177, G0409, G0463, G0512, G0560, H0002, H0004, H0031, H0034, H0036, H0037, H0039, H0040, H2000, H2010, H2011, H2013-H2020, T1015
<b>Partial Hospitalization or Intensive Outpatient</b> *must be with mental health provider or include appropriate dx codes	<b>CPT:</b> 90791, 90792, 90832-90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99252-99255  <b>+ must include POS:</b> 52



## (FUA) Follow-Up After Emergency Department Visit for Substance Use (continued)

<b>Non-residential substance abuse treatment facility visit</b> *must be with mental health provider or include appropriate dx codes	<b>CPT:</b> 90791, 90792, 90832-90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99252-99255  <b>+ must include POS:</b> 57 or 58
<b>Community Mental Health Center visit</b> *must be with mental health provider or include appropriate dx codes	<b>CPT:</b> 90791, 90792, 90832-90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99252-99255  <b>+ must include POS:</b> 53
<b>Peer Support Service</b> *must include appropriate dx codes	<b>HCPCS:</b> G0140, G0177, H0024, H0025, H0038-H0040, H0046, H2014, H2023, S9445, T1012, T1016, T1017
<b>Opioid Treatment Service</b> *must include appropriate dx codes	<u>Weekly Treatment</u> <b>HCPCS:</b> G2074, G2075, G2076, G2077, G2080  <u>Monthly Treatment</u> <b>HCPCS:</b> G2069, G2086, G2087
<b>Substance Use Disorder Service</b>	<b>CPT:</b> 99408, 99409 <b>HCPCS:</b> G0396, G0397, G0443, H0001, H0005, H0007, H0015, H0015, H0022, H0047, H0050, H2035, H2036, T1006, T1012
<b>Substance Use Disorder Counselling and Surveillance</b>	<b>ICD10CM:</b> Z71.41, Z71.51
<b>Behavioral Health Screening or Assessment for SUD or mental health disorders</b>	<b>CPT:</b> 99408, 99409 <b>HCPCS:</b> G0396, G0397, G0442, G2011, H0001, H0002, H0031, H0049

(continued)



## (FUA) Follow-Up After Emergency Department Visit for Substance Use (continued)

Substance Use Service	<b>HCPCS:</b> H0006, H0028
Diagnosis of SUD (SUD)	<b>ICD10CM:</b> F10.10, F10.120, F10.121, F10.129, F10.130, F10.131, F10.132, F10.139, F10.14, F10.150, F10.151, F10.159, F10.180, F10.181, F10.182, F10.188 *list does not include all due to amount
Substance Use	<b>ICD10CM:</b> F10.90, F10.920, F10.921, F10.929, F10.930, F10.931, F10.932, F10.939, F10.94, F10.950, F10.951, F10.959 *list does not include all due to amount
Drug Overdose	<b>ICD10CM:</b> T40.OX1A, T40.OX1D, T40.OX1S, T40.OX4A *list does not include all due to amount
Hospice Care	<b>HCPCS:</b> G9473-G9479, Q5003-Q5008, Q5010, S9126, T2042-T2046, G0182 <b>CPT:</b> 99377, 99378

\*Codes subject to change

### Action

- ✓ Complete one of the following within seven days of ED visit:
  - The following must be completed either with a mental health provider OR any provider if diagnosis of SUD, substance use or drug overdose is included
    - Outpatient visit, telehealth visit, telephone visit, e-visit or virtual check-in
    - Intensive outpatient encounter or partial hospitalization
    - Non-residential substance abuse treatment facility visit
    - Community mental health center visit
  - Peer support service with any diagnosis of SUD, substance use or drug overdose
  - Opioid treatment service that bills monthly or weekly with any diagnosis of SUD, substance use or drug overdose

(continued)



# (FUA) Follow-Up After Emergency Department Visit for Substance Use (continued)

- Substance use disorder service
- Substance use disorder counselling and surveillance
- Behavioral health screening or assessment for SUD or mental health disorders
- Substance use service
- Appropriate medications dispensed

✓ If the follow up does not occur within the first seven days of ED visit, please schedule the appointment to occur within 30 days

## Tips and Best Practices

- ✓ Schedule the seven-day follow-up visit within five days to allow flexibility in rescheduling
- ✓ Follow-up visit can occur on same day of ED visit
- ✓ Follow-up visit does not need to be with a mental health provider if diagnosis of SUD, substance use or drug overdose is included on claim
- ✓ Refer member to a provider who specializes in treating substance use or substance-induced disorders
- ✓ Review when to schedule office appointments versus seek urgent care and emergency department care

## Required Exclusions

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above or medical record documentation:

- ✓ Members who use hospice services any time during the measurement year
- ✓ Members who die anytime during the measurement year



# (FUI) Follow-Up After High-Intensity Care for Substance Use Disorder

## Product Lines:

- ✓ WellCare (Medicare)
- ✓ MI Coordinated Health (Dual Eligible Plan)

## Description

Measure evaluates percentage of acute inpatient hospitalizations, residential treatment or withdrawal management visits for a diagnosis of substance use disorder among members 13 years of age and older that result in a follow-up visit or service for substance use disorder

Two rates are reported:

- ✓ Visits or discharges for which the member received follow-up for substance use disorder within 30 days after the visit or discharge
- ✓ Visits or discharges for which the member received follow-up for substance use disorder within 7 days after the visit or discharge

Description	Codes*
<b>Outpatient Visit</b> *must include appropriate dx codes	<b>CPT:</b> 90791, 90792, 90832-90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99252-99255  <b>Include Outpatient POS code from following:</b> POS: 03, 05, 07, 09, 11-20, 22, 33, 49, 50, 71, 72

(continued)



## (FUI) Follow-Up After High-Intensity Care for Substance Use Disorder (continued)

<b>BH Outpatient Visit</b> *must include appropriate dx codes	<b>CPT:</b> 98000-98007, 98960-98962, 99078, 99202-99205, 99211-99215, 99242-99245, 99341-99345, 99347-99350, 99381-99387, 99391-99397, 99401-99404, 99411, 99412, 99483, 99492-99494, 99510 <b>HCPCS:</b> G0155, G0176, G0177, G0409, G0463, G0512, G0560, H0002, H0004, H0031, H0034, H0036, H0037, H0039, H0040, H2000, H2010, H2011, H2013-H2020, T1015
<b>Partial Hospitalization or Intensive Outpatient</b> *must include appropriate dx codes	<b>CPT:</b> 90791, 90792, 90832-90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99252-99255  <b>+ must include POS:</b> 52
<b>Non-residential substance abuse treatment facility visit</b> *must be with mental health provider or include appropriate dx codes	<b>CPT:</b> 90791, 90792, 90832-90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99252-99255  <b>+ must include POS:</b> 57 or 58
<b>Community Mental Health Center visit</b> *must be with mental health provider or include appropriate dx codes	<b>CPT:</b> 90791, 90792, 90832-90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99252-99255  <b>+ must include POS:</b> 53
<b>Peer Support Service</b> *must include appropriate dx codes	<b>HCPCS:</b> G0140, G0177, H0024, H0025, H0038-H0040, H0046, H2014, H2023, S9445, T1012, T1016, T1017

(continued)



## (FUI) Follow-Up After High-Intensity Care for Substance Use Disorder (continued)

<b>Opioid Treatment Service</b> *must include appropriate dx codes	<u>Weekly Treatment</u> <b>HCPCS:</b> G2074, G2075, G2076, G2077, G2080  <u>Monthly Treatment</u> <b>HCPCS:</b> G2069, G2086, G2087
<b>Substance Use Disorder Service</b>	<b>CPT:</b> 99408, 99409 <b>HCPCS:</b> G0396, G0397, G0443, H0001, H0005, H0007, H0015, H0015, H0022, H0047, H0050, H2035, H2036, T1006, T1012
<b>Substance Use Disorder Counselling and Surveillance</b>	<b>ICD10CM:</b> Z71.41, Z71.51
<b>Substance Use Service</b>	<b>HCPCS:</b> H0006, H0028
<b>Diagnosis of SUD (SUD)</b>	<b>ICD10CM:</b> F10.10, F10.120, F10.121, F10.129, F10.130, F10.131, F10.132, F10.139, F10.14, F10.150, F10.151, F10.159, F10.180, F10.181, F10.182, F10.188 *list does not include all due to amount
<b>Hospice Care</b>	<b>HCPCS:</b> G9473-G9479, Q5003-Q5008, Q5010, S9126, T2042-T2046, G0182 <b>CPT:</b> 99377, 99378

\*Codes subject to change

### Action

- ✓ Complete one of the following within seven days of visit or discharge (all MUST include diagnosis of substance use disorder):
  - Outpatient visit, telehealth visit, telephone visit, e-visit or virtual check-in
  - Intensive outpatient encounter or partial hospitalization
  - Non-residential substance abuse treatment facility visit
  - Community mental health center visit
  - Substance use disorder service



## (FUI) Follow-Up After High-Intensity Care for Substance Use Disorder (continued)

- Substance use disorder counselling and surveillance
- Opioid treatment service that bills monthly or weekly
- Residential behavioral health treatment
- Peer support services
- Appropriate medications dispensed

- ✓ If the follow up does not occur within the first seven days of visit or discharge, please schedule the appointment to occur within 30 days

### Tips and Best Practices

- ✓ Schedule the seven-day follow-up visit within five days to allow flexibility in rescheduling
- ✓ Follow-up visit can occur on same day of visit or discharge
- ✓ Follow-up visit does not need to be with any specific provider type
- ✓ Review when to schedule office appointments versus seek urgent care and emergency department care

### Required Exclusions

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above or medical record documentation:

- ✓ Members who use hospice services any time during the measurement year
- ✓ Members who die anytime during the measurement year



# (FUH) Follow-Up After Hospitalization for Mental Illness

## Product Lines

- ✓ Meridian (Medicaid)
- ✓ MI Coordinated Health (Dual Eligible Plan)
- ✓ Ambetter from Meridian (Marketplace)
- ✓ WellCare (Medicare)

## Description

Measure evaluates percentage of discharges for members 6 years of age and older who were hospitalized for a principal diagnosis of mental illness, or any diagnosis of intentional self-harm, and had a mental health follow-up service.

### Two rates are reported:

- 1 Discharges for which the member received follow-up within 30 days of discharge
- 2 Discharges for which the member received follow-up within 7 days of discharge

Description	Codes*
<b>BH Outpatient Visit</b> (must be with a mental health provider or include a diagnosis of mental health disorder)	<b>CPT:</b> 98000-98007, 98960-98962, 99078, 99202-99205, 99211-99215, 99242-99245, 99341-99345, 99347-99350, 99381-99387, 99391-99397, 99401-99404, 99411, 99412, 99483, 99492-99494, 99510 <b>HCPCS:</b> G0155, G0176, G0177, G0409, G0463, G0560, H0002, H0004, H0031, H0034, H0036, H0037, H0039, H0040, H2000, H2010, H2011, H2013-H2020, T1015

(continued)



## (FUH) Follow-Up After Hospitalization for Mental Illness (continued)

Description	Codes*
<b>Outpatient Visit with Unspecified Setting</b> (must also include outpatient POS code <b>and</b> be with a mental health provider or include a diagnosis of mental health disorder)	<b>CPT:</b> 90791, 90792, 90832-90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99252-99255 <b>AND</b> <b>Outpatient POS:</b> 03, 05, 07, 09, 11-20, 22, 33, 49, 50, 52, 53, 71, 72
<b>Telephone Visits</b>	<b>CPT:</b> 98966-98968, 99441-99443
<b>Partial Hospitalization or Intensive Outpatient</b>	<b>HCPCS:</b> G0410, G0411, H0035, H2001, H2012, S0201, S9480, S9484, S9485
<b>Psychiatric Collaborative Care Management</b>	<b>CPT:</b> 99492-99494 <b>HCPS:</b> G0512
<b>Transitional Care Management</b>	<b>CPT:</b> 99495, 99496
<b>Peer Support Services</b>	<b>HCPCS:</b> G0140, G0177, H0024, H0025, H0038, H0039, H0040, H0046, H2014, H2023, S9445, T1012, T1016
<b>Psychiatric Residential Treatment</b>	<b>HCPS:</b> H0017, H0018, H0019, T2048
<b>Electroconvulsive Therapy</b>	<b>CPT:</b> 90870
<b>Hospice Care</b>	<b>HCPCS:</b> G9473-G9479, Q5003-Q5008, Q5010, S9126, T2042-T2046, G0182 <b>CPT:</b> 99377, 99378

\*Codes subject to change

### Action

✓ Complete one of the following within seven days post-discharge:

- Outpatient visit with a mental health provider
- Outpatient visit with any diagnosis of mental health disorder
- Telehealth or telephone visit with a mental health provider or diagnosis of mental health disorder
- Transitional care management services with a mental health provider or diagnosis of mental health disorder



## (FUH) Follow-Up After Hospitalization for Mental Illness (continued)

- Intensive outpatient encounter or partial hospitalization
- Community mental health center visit
- Psychiatric collaborative care management
- Peer support services with a diagnosis of mental health disorder
- Psychiatric residential treatment
- Electroconvulsive therapy

✓ If the follow-up does not occur within the first seven days post-discharge, please schedule the appointment to occur within 30 days

### Tips and Best Practices

- ✓ Schedule the seven-day follow up visit within five days to allow flexibility in rescheduling. Involve the patient's caregiver regarding the follow-up plan after inpatient discharge
- ✓ Follow-up does not need to be with a mental health provider, if there is a diagnosis of mental health disorder included
- ✓ Schedule follow-up appointments prior to discharge and include the date and time on discharge instructions
- ✓ Follow-up visit should not be completed on the date of discharge
- ✓ Contact Meridian for assistance locating a mental health provider
- ✓ Identify barriers for the patient and address their concerns
- ✓ Review when to schedule office appointments versus seek urgent care and emergency department care

### Required Exclusions

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above or medical record documentation:

- ✓ Members who use hospice services any time during the measurement year
- ✓ Members who die any time during the measurement year



# (FUM) Follow-Up After Emergency Department Visit for Mental Illness

## Product Lines:

- ✓ Meridian (Medicaid)
- ✓ MI Coordinated Health (Dual Eligible Plan)
- ✓ Ambetter from Meridian (Marketplace)
- ✓ WellCare (Medicare)

## Description

Measure evaluates percentage of emergency department (ED) visits for members 6 years of age and older with a principal diagnosis of mental illness, or any diagnosis of intentional self-harm, and had a mental health follow-up service.

Two rates are reported:

- ✓ ED visits for which the member received follow-up within 30 days of discharge
- ✓ ED visits for which the member received follow-up within 7 days of discharge

Description	Codes*
BH Outpatient Visit	<b>CPT:</b> 98960–98962, 99078, 99202–99205, 99211–99215, 99242–99245, 99341–99345, 99347–99350, 99381–99387, 99391–99397, 99401–99404, 99411, 99412, 99483, 99492–99494, 99510 <b>HCPGs:</b> G0155, G0176, G0177, G0409, G0463, G0512, H0002, H0004, H0031, H0034, H0036, H0037, H0039, H0040, H2000, H2010, H2011, H2013–H2020, T1015

(continued)



## (FUM) Follow-Up After Emergency Department Visit for Mental Illness (continued)

Description	Codes*
Outpatient Visit with Unspecified Setting	<b>CPT:</b> 90791, 90792, 90832-90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99252-99255 <b>AND</b>
	<b>POS:</b> 03, 05, 07, 09, 11-20, 22, 33, 49, 50, 52, 53, 71, 72
Telephone Visits	<b>CPT:</b> 98966-98968, 99441-99443
Partial Hospitalization or Intensive Outpatient	<b>HCPCS:</b> G0410, G0411, H0035, H2001, H2012, S0201, S9480, S9484, S9485
Psychiatric Collaborative Care Management	<b>CPT:</b> 99492-99494 <b>HCPCS:</b> G0512
Transitional Care Management	<b>CPT:</b> 99495, 99496
Peer Support Services	<b>HCPCS:</b> G0140, G0177, H0024, H0025, H0038, H0039, H0040, H0046, H2014, H2023, S9445, T1012, T1016
Psychiatric Residential Treatment	<b>HCPCS:</b> H0017, H0018, H0019, T2048
Electroconvulsive Therapy	<b>CPT:</b> 90870
Hospice Care	<b>HCPCS:</b> G9473-G9479, Q5003-Q5008, Q5010, S9126, T2042-T2046, G0182 <b>CPT:</b> 99377, 99378

\*Codes subject to change

### Action

✓ Complete one of the following with seven days of ED visit:

- Outpatient visit with any diagnosis of mental health disorder
- Telehealth or telephone visit with a diagnosis of mental health disorder
- Transitional care management services with a diagnosis of mental health disorder



## (FUM) Follow-Up After Emergency Department Visit for Mental Illness (continued)

- Intensive outpatient encounter or partial hospitalization
- Community mental health center visit
- Psychiatric collaborative care management
- Peer support services with a diagnosis of mental health disorder
- Psychiatric residential treatment
- Electroconvulsive therapy

✓ If the follow up does not occur within the first seven days of ED visit, please schedule the appointment to occur within 30 days

### Tips and Best Practices

- Schedule the seven-day follow up visit within five days to allow flexibility in rescheduling. Involve the patient's caregiver regarding the follow-up plan
- Follow-up visit does not need to be with a mental health provider
- Follow-up visit can occur on same day of ED visit
- Identify barriers for the patient and address their concerns
- Review when to schedule office appointments versus seek urgent care and emergency department care

### Required Exclusions

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above or medical record documentation:

- Members who use hospice services any time during the measurement year
- Members who die anytime during the measurement year



# (IET) Initiation and Engagement of Substance Use Disorder Treatment

## Product Lines

- ✓ Meridian (Medicaid)
- ✓ MI Coordinated Health (Dual Eligible Plan)
- ✓ Ambetter from Meridian (Marketplace)
- ✓ WellCare (Medicare)

## Description

Measure evaluates percentage of new substance use disorder (SUD) episodes that result in treatment initiation and engagement.

### Two rates are reported:

- 1 **Initiation of SUD Treatment:** percentage of new SUD episodes that result in treatment initiation through an inpatient SUD admission, outpatient visit, intensive outpatient encounter or partial hospitalization, telehealth, or medication treatment within 14 days
- 2 **Engagement of SUD Treatment:** percentage of new SUD episodes that have evidence of treatment engagement within 34 days of initiation

Description	Prescription
<b>Outpatient Visit with Unspecified Setting</b>	<b>CPT:</b> 90791, 90792, 90832-90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99252-99255 <b>AND</b> <b>POS:</b> 03, 05, 07, 09, 11-20, 22, 33, 49, 50, 52, 53, 56, 57, 58 71, 72

(continued)



## (IET) Initiation and Engagement of Substance Use Disorder Treatment (continued)

<b>BH Outpatient Visit</b>	<b>CPT:</b> 98960-98962, 99078, 99202-99205, 99211-99215, 99242-99245, 99341-99345, 99347-99350, 99381-99387, 99391-99397, 99401-99404, 99411, 99412, 99483, 99492-99494, 99510 <b>HCPCS:</b> G0155, G0176, G0177, G0409, G0463, G0512, H0002, H0004, H0031, H0034, H0036, H0037, H0039, H0040, H2000, H2010, H2011, H2013-H2020, T1015
<b>Intensive Outpatient Encounter or Partial Hospitalization</b>	<b>HCPCS:</b> G0410, G0411, H0035, H2001, H2012, S0201, S9480, S9484, S9485
<b>Substance Use Disorder Counseling and Surveillance</b>	<b>ICD10CM:</b> Z71.41, Z71.51
<b>Weekly or Monthly Opioid Treatment Service</b>	<b>HCPCS:</b> G2071, G2074-G2077, G2080, G2067-G2070, G2072, G2073, G2086, G2087
<b>Telehealth Visit</b>	<b>POS:</b> 02, 10
<b>Telephone Visit</b>	<b>CPT:</b> 98966-98968, 99441-99443
<b>E-Visit or Virtual Check-In</b>	<b>CPT:</b> 98970-98972, 98980, 98981, 99421-99423, 99457, 99458 <b>HCPCS:</b> G0071, G2010, G2012, G2250-G2252
<b>Hospice Care</b>	<b>HCPCS:</b> G9473-G9479, Q5003-Q5008, Q5010, S9126, T2042-T2046, G0182 <b>CPT:</b> 99377, 99378

\*Codes subject to change



## (IET) Initiation and Engagement of Substance Use Disorder Treatment (continued)

### Action

- ✓ Complete one of the following within 14 days (must include SUD diagnosis):
  - Outpatient visit, telehealth, telephone visit or virtual check-in
  - Intensive outpatient encounter or partial hospitalization
  - Non-residential substance abuse treatment facility visit
  - Community mental health center visit
  - Substance use disorder counseling and surveillance
  - Weekly or monthly opioid treatment service
  - Dispense medication to treat alcohol use disorder or opioid use disorder
- ✓ Complete one of the above options again within 34 days of first treatment

### Tips and Best Practices

- ✓ Schedule the initial 14-day treatment visit within 10 days of new SUD episode to allow flexibility in rescheduling
- ✓ At the end of the initial appointment, schedule two more appointments to occur within 34 days of the initial appointment
- ✓ Offer SUD treatment option on-site or support treatment scheduling with accessible in-network treatment facility during patient visit and have referrals easily accessible
- ✓ When treating a patient for issues related to SUD, code that diagnosis on every claim
- ✓ Contact Meridian for assistance locating SUD treatment provider, if needed
- ✓ Not all medications may be covered. For a list of covered medications, refer to the drug formulary at **mimeridian.com** or **www.wellcare.com**

(continued)



## (IET) Initiation and Engagement of Substance Use Disorder Treatment (continued)

### Required Exclusions

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above or medical record documentation:

- ✓ Members who use hospice services any time during the measurement year
- ✓ Members who die any time during the measurement year
- ✓ Members who were treated for SUD within the previous 194 days

\*For the follow-up treatments, include an ICD-10 diagnosis for Alcohol or Other Drug Dependence from the Mental, Behavioral and Neurodevelopmental Disorder Section of ICD-10 along with a procedure code for the preventive service, evaluation, and management consultation or counseling service.

.....



## (MI2.6) Timely Transmission of Care Transition Record to Health Care Professional

### Product Lines:

- ✓ MI Coordinated Health (Dual Eligible Plan)

### Description

Measure evaluates the percentage of members for whom a transition record was transmitted to the facility, primary physician, or other health professional designated for follow-up care on the day of discharge through two days after discharge

(continued)



## (MI2.6) Timely Transmission of Care Transition Record to Health Care Professional (continued)

### Action

- ✓ Discharging facility should send primary care or ongoing care provider a copy of the patient's discharge summary within two days of discharge

### Tips and Best Practices

- ✓ Investigate and implement a process to receive auto alerts when a patient is discharged from an inpatient facility
- ✓ Discharging facility should date and time stamp when the discharge summary was sent to outpatient care provider
- ✓ Outpatient provider should date and time stamp when the discharge summary was received in the member's medical record

### Required Exclusions

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table below or medical record documentation:

- ✓ Members who died during the inpatient stay or during hospice care
- ✓ Members who left against medical advice or discontinued care

Description	Codes*
Left against Medical Advice or Discontinued Care	Discharge Status: 07
Member Expired	Discharge Status: 20, 40, 41, 42

\*Codes subject to change



## (MI5.6) Medication Review

### Product Lines:

- ✓ MI Coordinated Health (Dual Eligible Plan)

### Description

Measure evaluates the percentage of members who had at least one medication review conducted by a prescribing practitioner or clinical pharmacist during the measurement year and the presence of a medication list in the medical record.

Description	Codes*
Medication Review (codes must be submitted for both Medication List and Medication Review for credit)	<b>Medication List-</b> <b>CPT-CAT-II:</b> 1159F <b>HCPCS:</b> G8427  <b>Medication Review-</b> <b>CPT-CAT-II:</b> 1160F <b>CPT:</b> 90863, 99605, 99606, 99483
Transitional Care Management Services	<b>CPT:</b> 99495, 99496
Hospice Care	<b>HCPCS:</b> G9473-G9479, Q5003-Q5008, Q5010, S9126, T2042-T2046, G0182 <b>CPT:</b> 99377, 99378

\*Codes subject to change

### Action

- ✓ Perform an annual review of the patient's medications, including prescription medications, over-the-counter medications and herbal or supplemental therapies. This must be completed by a prescribing provider or a clinical pharmacist.

(continued)



## (MI5.6) Medication Review *(continued)*

### Tips and Best Practices

- ✓ If member is not taking any medication, document this in the record to count towards this measure
- ✓ Medication review cannot come from an acute inpatient setting, but is acceptable from an emergency department visit
- ✓ Medication review must be comprehensive, not limited to side effects for a single medication at the time of prescription

### Required Exclusions

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above or medical record documentation:

- ✓ Members who use hospice services any time during the measurement year
- ✓ Members who die any time during the measurement year

---



## (MI7.3) Annual Dental Visit/Adults: Any Dental Visit

### Product Lines:

- ✓ MI Coordinated Health (Dual Eligible Plan)
- ✓ Meridian (Medicaid)

### Description

Measure evaluates the percentage of members 19 and older who had one or more dental visits with a dental provider during the measurement year.

*(continued)*



## (MI7.3) Annual Dental Visit/Adults: Any Dental Visit *(continued)*

DESCRIPTION	CODES*
Diagnostic Care	<b>CDT:</b> D0120, D0140, D0145, D150, D5160, D0170, D0171, D0171, D0180, D0190, D0191, D0210, D0220, D0230, D0240, D0240, D0250, D0260, D0270, D0272-D0274, D0277, D0290, D0310, D0320-D0322, D0330, D0340, D0350-D0351, D0364-D0369, D0370-D0371, D0380-D0386, D0391, D0393-D0395, D0415-D0418, D0421, D0425, D0431, D0460, D0470, D0472-D0479, D0480-D0486, D0502, D0601-D0603, D0999  <b>CPT:</b> 70300, 70310, 70320, 70350, 70355
Preventive Care	<b>CDT:</b> D1110, D1206, D1208, D1310, D1320, D1351-D1353, D1510, D1515, D1520, D1525, D1550, D1555, D1999
Restorative Care	<b>CDT:</b> D2140, D2150, D2160-D2161, D2330-D2332, D2335, D2390-D2394, D2410, D2420D, 2430D, 25110, D2520, D2530, D2542-D2544, D2620, D2630, D2642-D2644, D2650-D2652, D2662-D2664, D2710, D2720-D2722, D2740, D2750-D2752, D2780-D2783, D2790-D2794, D2799, D2910, D2915, D2920-D2921, D2929, D2930-D2934, D2940-D2941, D2949, D2950-2955, D2957, D2960-D2962, D2970-D2971, D2975, D2980-D2983, D2990, D2999

*(continued)*



## (MI7.3) Annual Dental Visit/Adults: Any Dental Visit *(continued)*

Endodontics	<b>CDT:</b> D3110, D3120-D3222, D3230, D3240, D3310, D3320, D3330-D3333, D3346-D3348, D3351-D3353, D3355-D3357, D3410, D3421, D3425-D3429, D3430-D3432, D3450, D3460, D3470, D3910, D3920, D3950, D3999
Periodontics	<b>CDT:</b> D4210-D4212, D4230-D4231, D4240-D4241, D4245, D4249, D4260-D4268, D4270, D4273-D4278, D4320-D4321, D4341, D4342, D4355, D4381, D4910, D4920-D4921, D4999
Prosthodontics	<b>CDT:</b> D5110, D5120, D5130, D5140, D5211-D5214, D5225-D5226, D5281, D5410-D5411, D5421-D5422, D5510, D5520, D5610, D5620, D5630, D5640, D5650, D5660, D5670-D5671, D5710-D5711, D5720-D5721, D5730-D5731, D5740-D5741, D5750-D5751, D5760-D5761, D5810-D5811, D5820-D5821, D5850-D5851, D5862-D5867, D5875, D5899, D5994
Implant Services	<b>CDT:</b> D6000-D6199
Prosthodontics, Fixed	<b>CDT:</b> D6200-D5999
Oral & maxillofacial Surgery	<b>CDT:</b> D7000-D7999
Orthodontics	<b>CDT:</b> D8000-D8999
Adjunctive General Services	<b>CDT:</b> D9000-D9999
Hospice Care	<b>HCPSCS:</b> G9473-G9479, Q5003-Q5008, Q5010, S9126, T2042-T2046, G0182, <b>CPT:</b> 99377, 99378

\*Codes subject to change

*(continued)*



## (MI7.3) Annual Dental Visit/Adults: Any Dental Visit (continued)

### Action

- ✓ Primary care providers should encourage patients to schedule annual dental visits for oral health care and routine checkups
- ✓ Dental practitioner should perform dental visit at least one per year to prevent disease, reverse disease processes, prevent progression of caries, and reduce incidence of future lesions
- ✓ Dental practitioner is defined as:
  - Doctor of Dental Surgery (DDS)
  - Doctor of Dental Medicine (DMD)
  - Certified and Licensed Dental Hygienist

### Tips and Best Practices

- ✓ Any visit with a dental practitioner during the measurement year is acceptable, whether preventive or not
- ✓ Encourage new patients to establish a dental home to ensure good routine oral healthcare and follow ups
- ✓ Send reminders every six months to schedule for periodic exams and prophylaxis (cleanings)

### Required Exclusions

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above or medical record documentation:

- ✓ Members who use hospice services any time during the measurement year

---



# (URI) Appropriate Treatment for Upper Respiratory Infection

## Product Lines:

- ✓ Meridian (Medicaid)
- ✓ Ambetter from Meridian (Marketplace)
- ✓ WellCare (Medicare)
- ✓ MI Coordinated Health (Dual Eligible Plan)

## Description

Measure evaluates the percentage of episodes for members 3 months of age and older with a diagnosis of upper respiratory infection (URI) that did not result in an antibiotic dispensing event.

Description	Codes*
Hospice Care	<b>HCPCS:</b> G9473-G9479, Q5003-Q5008, Q5010, S9126, T2042-T2046, G0182 <b>CPT:</b> 99377, 99378

\*Codes subject to change

## Action

- ✓ Evaluate each patient with upper respiratory infection to determine if antibiotics are needed. Avoid inappropriately prescribing, which can lead to antibiotic resistance

## Tips and Best Practices

- ✓ Provide handouts explaining that viruses, not bacteria, cause colds and flu. CDC resource found here: [www.cdc.gov/antibiotic-use/week/toolkit.html#anchor\\_1538595518](http://www.cdc.gov/antibiotic-use/week/toolkit.html#anchor_1538595518)
- ✓ If patients have comorbid or competing diagnoses, submit claims for these diagnoses to remove patient from the HEDIS measure. Most common competing and comorbid diagnoses are:
  - COPD
  - Emphysema
  - Sickle cell disease
  - Cystic fibrosis



## (URI) Appropriate Treatment for Upper Respiratory Infection *(continued)*

- Acute pharyngitis
- Acute tonsillitis
- Acute or chronic sinusitis
- Pneumonia
- UTI

### Required Exclusions

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above or medical record documentation:

- ✓ Members who use hospice services any time during the measurement year
- ✓ Members who die any time during the measurement year

---



## (SNS-E) Social Need Screening and Intervention

### Product Lines:

- ✓ Meridian (Medicaid)
- ✓ MI Coordinated Health (Dual Eligible Plan)
- ✓ Ambetter from Meridian (Marketplace)
- ✓ WellCare (Medicare)

### Description

Measure evaluates percentage of members who were screened, using prespecified instruments, at least once during the measurement year for unmet food, housing and transportation needs, and received a corresponding intervention if they screened positive.



## (SNS-E) Social Need Screening and Intervention (continued)

- **Food Screening:** The percentage of members who were screened for food insecurity between 1/1/MY-12/1/MY
- **Food Intervention:** Of those that screened positive for food insecurity, the percentage who received a corresponding intervention within 1 month (31 days total) of screening positive for food insecurity
- **Housing Screening:** The percentage of members who were screened for housing instability, homelessness or housing inadequacy between 1/1/MY-12/1/MY
- **Housing Intervention:** Of those that screened positive for housing instability, homelessness or housing inadequacy, the percentage who received a corresponding intervention within 1 month (31 days total)
- **Transportation Screening:** The percentage of members who were screened for transportation insecurity between 1/1/MY-12/1/MY
- **Transportation Intervention:** Of those that screened positive for transportation insecurity, the percentage who received a corresponding intervention within 1 month (31 days total) of screening positive for transportation insecurity

### Action

- ✓ During each visit, use an approved screening instrument (listed in table below) to ask your patients about their social needs related to:
  - **Food insecurity:** Uncertain, limited, or unstable access to food that is: adequate in quantity and in nutritional quality; culturally acceptable; safe and acquired in socially acceptable ways

(continued)



## (SNS-E) Social Need Screening and Intervention *(continued)*

- **Housing instability:** Currently consistently housed but experiencing any of the following circumstances in the past 12 months: being behind on rent or mortgage, multiple moves, cost burden or risk of eviction
- **Homelessness:** Currently living in an environment that is not meant for permanent human habitation (such as cars, parks, abandoned buildings), not having a consistent place to sleep at night, or because of economic difficulties, currently living in a shelter, motel temporary or transitional living situation
- **Housing Inadequacy:** Housing does not meet standard habitability standards
- **Transportation insecurity:** Uncertain, limited or no access to safe, reliable, accessible, affordable, and socially acceptable transportation infrastructure and modalities necessary for maintaining one's health, well-being or livelihood

✓ If your patient screens positive for any social needs, complete one of the following interventions: assistance, assessment, counseling, coordination, education, evaluation of eligibility, provision or referral. Complete this in the same visit that the need is identified, if possible. Applicable codes are listed in the table below:



## (SNS-E) Social Need Screening and Intervention (continued)

Description	Codes* Instrument	Screening Item LOINC Codes	Positive Finding LOINC Codes
Food Insecurity Instruments	Accountable Health Communities (AHC) Health-Related Social Needs (HRSN) Screening Tool	88122-7 88123-5	LA28397-0 LA6729-3 LA28397-0 LA6729-3
	American Academy of Family Physicians (AAFP) Social Needs Screening Tool	88122-7	LA28397-0 LA6729-3
		88123-5	LA28397-0 LA6729-3
	American Academy of Family Physicians (AAFP) Social Needs Screening Tool—short form	88122-7	LA28397-0 LA6729-3
		88123-5	LA28397-0 LA6729-3
	Health Leads Screening Panel®1	95251-5	LA33-6
	Hunger Vital Sign™1 (HVS)	88124-3	LA19952-3
	Protocol for Responding to and Assessing Patients' Assets, Risks and Experiences [PRAPARE]®1	93031-3	LA30125-1
	Safe Environment for Every Kid (SEEK)®1	95400-8	LA33-6
		95399-2	LA33-6
	U.S. Household Food Security Survey [U.S. FSS]	95264-8	LA30985-8 LA30986-6
	U.S. Adult Food Security Survey [U.S. FSS]	95264-8	LA30985-8 LA30986-6
	U.S. Child Food Security Survey [U.S. FSS]	95264-8	LA30985-8 LA30986-6
	U.S. Household Food Security Survey-Six-Item Short Form [U.S. FSS]	95264-8	LA30985-8 LA30986-6
	We Care Survey	96434-6	LA32-8



## (SNS-E) Social Need Screening and Intervention (continued)

Food Insecurity Interventions	<p><b>CPT:</b> 96156, 96160, 96161, 97802-97804</p> <p><b>HCPCS:</b></p> <p><b>S5170:</b> Home delivered meals, including preparation; per meal</p> <p><b>S9470:</b> Nutritional counseling, dietitian visit</p> <p><b>SNOMED CT:</b></p> <p><b>1759002:</b> Assessment of nutritional status</p> <p><b>61310001:</b> Nutrition education</p> <p><b>103699006:</b> Referral to dietitian</p> <p><b>30844001:</b> Referral to social worker</p> <p><b>385767005:</b> Meals on wheels provision education</p> <p><b>710925007:</b> Provision of food</p> <p><b>464691000124107:</b> Counseling for barriers to achieving food security</p> <p><b>713109004:</b> Referral to community meals service</p> <p><b>1002223009:</b> Assessment of progress towards goals to achieve food security</p> <p><b>1230338004:</b> Referral to charitable organization</p> <p><b>441041000124100:</b> Counseling about nutrition</p> <p><b>1268662008:</b> Coordination of resources to address social needs</p> <p><b>711069006:</b> Coordination of care plan</p> <p><b>710824005:</b> Assessment of health and social care needs</p> <p><b>663211000124100:</b> Referral to peer support specialist</p> <p><b>663081000124100:</b> Referral to State Funded Food Assistance Program</p> <p><b>662151000124104:</b> Assistance with application for State Funded Food Assistance Program</p> <p><b>662651000124105:</b> Evaluation of eligibility for State Funded Food Assistance Program</p>
-------------------------------	---



## (SNS-E) Social Need Screening and Intervention (continued)

	Instrument	Screening Item LOINC Codes	Positive Finding LOINC Codes
Housing Instability, Homelessness, and Housing Inadequacy Instruments	Accountable Health Communities (AHC) Health-Related Social Needs (HRSN) Screening Tool	71802-3	LA31994-9 LA31995-6
	American Academy of Family Physicians (AAFP) Social Needs Screening Tool	99550-6	LA33-6
	American Academy of Family Physicians (AAFP) Social Needs Screening Tool—short form	71802-3	LA31994-9 LA31995-6
	Children's Health Watch Housing Stability Vital Signs™ <sup>1</sup>	98976-4	LA33-6
		98977-2	≤2
		98978-0	LA33-6
	Health Leads Screening Panel® <sup>1</sup>	99550-6	LA33-6
	Protocol for Responding to and Assessing Patients' Assets, Risks and Experiences [PRAPARE]® <sup>1</sup>	93033-9	LA33-6
		71802-3	LA30190-5
	We Care Survey	96441-1	LA33-6
	WellRx Questionnaire	93669-0	LA33-6
	Accountable Health Communities (AHC) Health-Related Social Needs (HRSN) Screening Tool	96778-6	LA31996-4 LA28580-1 LA31997-2 LA31998-0

(continued)



## (SNS-E) Social Need Screening and Intervention *(continued)*

Instrument	Screening Item LOINC Codes	Positive Finding LOINC Codes
American Academy of Family Physicians (AAFP) Social Needs Screening Tool	96778-6	LA32691-0 LA28580-1 LA32693-6 LA32694-4 LA32695-1 LA32696-9 LA32001-2
American Academy of Family Physicians (AAFP) Social Needs Screening Tool—short form	96778-6	LA31996-4 LA28580-1 LA31997-2 LA31998-0 LA31999-8 LA32000-4 LA32001-2
Norwalk Community Health Center Screening Tool [NCHC]	99134-9	LA33-6
	99135-6	LA31996-4 LA28580-1 LA31997-2 LA31998-0 LA31999-8 LA32000-4 LA32001-2



## (SNS-E) Social Need Screening and Intervention *(continued)*

<p>Housing instability, Homelessness, and Inadequate Housing Interventions</p>	<p><b>CPT:</b> 96156, 96160, 96161</p> <p><b>SNOMED CT:</b></p> <p><b>1148447008:</b> Assessment for housing insecurity</p> <p><b>1148812007:</b> Assessment of progress toward goals to achieve housing</p> <p><b>1148814008:</b> Assessment of goals to achieve housing security</p> <p><b>710824005:</b> Assessment of health and social care needs</p> <p><b>711069006:</b> Coordination of care plan</p> <p><b>30844001:</b> Referral to social worker</p> <p><b>1230338004:</b> Referral to charitable organization</p> <p><b>470231000124107:</b> Counseling for social determinant of health risk</p> <p><b>1148813002:</b> Assessment of barriers in inadequate housing care plan</p> <p><b>1148815009:</b> Assessment of goals to achieve adequate housing</p> <p><b>1162436000:</b> Referral to legal aid</p> <p>*list does not include all SNOMED codes, due to high volume</p>
--	---



## (SNS-E) Social Need Screening and Intervention (continued)

	Transportation Insecurity Instruments	Screening Item LOINC Codes	Positive Finding LOINC Codes
Transportation Insecurity Instruments	Accountable Health Communities (AHC) Health-Related Social Needs (HRSN) Screening Tool	93030-5	LA33-6
	American Academy of Family Physicians (AAFP) Social Needs Screening Tool	99594-4	LA33-6
	American Academy of Family Physicians (AAFP) Social Needs Screening Tool—short form	99594-4	LA33093-8 LA30134-3
	Comprehensive Universal Behavior Screen (CUBS)	89569-8	LA29232-8 LA29233-6 LA29234-4
	Health Leads Screening Panel®1	99553-0	LA33-6
	Inpatient Rehabilitation Facility—Patient Assessment Instrument (IRF-PAI)—version 4.0 [CMS Assessment]	101351-5	LA30133-5 LA30134-3
	Outcome and assessment information set (OASIS) form—version E—Discharge from Agency [CMS Assessment]	101351-5	LA30133-5 LA30134-3
	Outcome and assessment information set (OASIS) form—version E—Resumption of Care [CMS Assessment]	101351-5	LA30133-5 LA30134-3
	Outcome and assessment information set (OASIS) form—version E—Start of Care [CMS Assessment]	101351-5	LA30133-5 LA30134-3
	Protocol for Responding to and Assessing Patients' Assets, Risks and Experiences [PRAPARE]®1	93030-5	LA30133-5 LA30134-3
PROMIS®1		92358-1	LA30024-6 LA30026-1 LA30027-9
WellRx Questionnaire		93671-6	LA33-6



## (SNS-E) Social Need Screening and Intervention (continued)

Transportation Interventions	<p><b>CPT:</b> 96156, 96160, 96161</p> <p><b>SNOMED CT:</b></p> <p><b>710824005:</b> Assessment of health and social care needs</p> <p><b>711069006:</b> Coordination of care plan</p> <p><b>30844001:</b> Referral to social worker</p> <p><b>1230338004:</b> Referral to charitable organization</p> <p><b>470231000124107:</b> Counseling for social determinant of health risk</p> <p><b>611281000124107:</b> Counseling for readiness to achieve transportation security</p> <p><b>611291000124105:</b> Counseling for barriers to achieve transportation security</p> <p><b>228615008:</b> Provision of transport</p> <p><b>611031000124101:</b> Education about rideshare program</p> <p><b>61106100124105:</b> Education about public transportation voucher program</p> <p>*list does not include all SNOMED codes, due to high volume</p>
------------------------------	---

### Tips and Best Practices

- ✓ Bring up social needs at each visit. Patients may need to be prompted to discuss non-medical issues with their provider
- ✓ Explain the purpose of the screening with the patient and offer support and resources as appropriate
- ✓ Allow for adequate time, space and privacy for the patient to discuss their social needs
- ✓ Educate your staff on the importance of screening, documenting, and coding patient's social needs



## (SNS-E) Social Need Screening and Intervention *(continued)*

### **Required Exclusions**

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above or medical record documentation:

- ✓ Members who use hospice services any time during the measurement year
- ✓ Members who die anytime during the measurement year

.....

# Medication Adherence

---



## (PDC-DIA) Medication Adherence for Diabetes Medications

### Product Lines:

- ✓ WellCare (Medicare)
- ✓ MI Coordinated Health (Dual Eligible Plan)

### Description

Measure evaluates percentage of Medicare Part D members 18 years and older with a prescription for diabetes medication who fill their prescription often enough to cover 80% or more of the time they are supposed to be taking the medication.

Diabetes Medications	
Biguanide Drugs	GLP-1 Receptor Agonists
Sulfonylurea Drugs	Meglitinide Drugs
Thiazolidinedione Drugs	Sodium Glucose Cotransporter 2 (SGLT2) Inhibitors
DiPeptidyl Peptidase (DPP)-4 Inhibitors	

### Action

- ✓ Urge patients to fill their prescriptions regularly to encourage medication adherence
- ✓ Encourage medication adherence by providing 90-day prescriptions. Your patient may be eligible for prescription delivery by mail through Express Scripts

### Tips and Best Practices

- ✓ Talk with patients about why they're on diabetes medications and why it's important to take their medication as prescribed and get refills promptly



## (PDC-DIA) Medication Adherence for Diabetes Medications *(continued)*

- ✓ Encourage patients to use a pillbox to organize their medications and to set an alarm on their smartphone or clock to remind them to take their medications
- ✓ Encourage patients to sign up for refill reminders at their pharmacy
- ✓ Monitor with scheduled follow-up appointments

### Required Exclusions

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above or medical record documentation:

- ✓ Members who use hospice services any time during the measurement year
- ✓ Members who die any time during the measurement year
- ✓ Members with a diagnosis of ESRD or dialysis coverage dates
- ✓ Members with one or more prescriptions for insulin

---



## (PDC-RAS) Medication Adherence for Hypertension - RAS antagonists

### Product Lines:

- ✓ WellCare (Medicare)
- ✓ MI Coordinated Health (Dual Eligible Plan)

### Description

Measure evaluates percentage of Medicare Part D members 18 years and older with a prescription for a blood pressure medication who fill their prescription often enough to cover 80% or more of the time they are supposed to be taking the medication.

*(continued)*



## (PDC-RAS) Medication Adherence for Hypertension - RAS antagonists (continued)

Hypertension Medications	
Angiotensin Converting Enzyme Inhibitors (ACEI)	Angiotensin Receptor Blockers (ARB)
Direct Renin Inhibitor Medications	

### Action

- ✓ Urge patients to fill their prescriptions regularly to encourage medication adherence
- ✓ Encourage medication adherence by providing 90-day prescriptions. Your patient may be eligible for prescription delivery by mail through Express Scripts

### Tips and Best Practices

- ✓ Talk with patients about why they're on hypertension medication and why it's important to take their medication as prescribed and get refills promptly
- ✓ Encourage patients to use a pillbox to organize their medications and to set an alarm on their smartphone or clock to remind them to take their medications
- ✓ Encourage patients to sign up for refill reminders at their pharmacy
- ✓ Monitor with scheduled follow-up appointments

### Required Exclusions

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above or medical record documentation:

- ✓ Members who use hospice services any time during the measurement year
- ✓ Members who die any time during the measurement year
- ✓ Members with a diagnosis of ESRD or dialysis coverage dates
- ✓ Members with one or more prescriptions for sacubitril/valsartan



## (PDC-STA) Medication Adherence for Cholesterol (Statins)

### Product Lines:

- ✓ WellCare (Medicare)
- ✓ MI Coordinated Health (Dual Eligible Plan)

### Description

Measure evaluates percentage of Medicare Part D members 18 years and older with a prescription for a cholesterol medication who fill their prescription often enough to cover 80% or more of the time they are supposed to be taking the medication.

### Action

- ✓ Urge patients to fill their prescriptions regularly to encourage medication adherence
- ✓ Encourage medication adherence by providing 90-day prescriptions. Your patient may be eligible for prescription delivery by mail through Express Scripts

### Tips and Best Practices

- ✓ Talk with patients about why they're on cholesterol medication and why it's important to take their medication as prescribed and get refills promptly
- ✓ Encourage patients to use a pillbox to organize their medications and to set an alarm on their smartphone or clock to remind them to take their medications
- ✓ Encourage patients to sign up for refill reminders at their pharmacy
- ✓ Monitor with scheduled follow-up appointments

(continued)



## (PDC-STA) Medication Adherence for Cholesterol (Statins) (continued)

### Required Exclusions

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above or medical record documentation:

- ✓ Members who use hospice services any time during the measurement year
- ✓ Members who die any time during the measurement year
- ✓ Members with a diagnosis of ESRD or dialysis coverage dates

---



## (SPC-E) Statin Therapy for Patients with Cardiovascular Disease

### Product Lines:

- ✓ Meridian (Medicaid)
- ✓ MI Coordinated Health (Dual Eligible Plan)
- ✓ WellCare (Medicare)

### Description

Measure evaluates percentage of males 21-75 years of age and females 40-75 years of age during the measurement year who were identified as having clinical atherosclerotic cardiovascular disease (ASCVD) and met the following criteria.

### Two rates are reported:

- ✓ **Received Statin Therapy:** Percentage of members who were dispensed at least one high-intensity or moderate intensity statin medication during the measurement year.



## (SPC-E) Statin Therapy for Patients with Cardiovascular Disease (continued)

✓ **Statin Adherence 80%:** Percentage of members who remained on a high-intensity or moderate-intensity statin medication for at least 80% of the treatment period.

Description	Prescription
<b>High-intensity statin therapy</b>	Atorvastatin 40-80 mg Amlodipine-atorvastatin 40-80 mg Rosuvastatin 20-40 mg Simvastatin 80 mg Ezetimibe-simvastatin 80 mg
<b>Moderate-intensity statin therapy</b>	Atorvastatin 10-20 mg Amlodipine-atorvastatin 10-20 mg Rosuvastatin 5-10 mg Simvastatin 20-40 mg Ezetimibe-simvastatin 20-40 mg Pravastatin 40-80 mg Lovastatin 40-60 mg Fluvastatin 40-80 mg Pitavastatin 1-4 mg

### Action

✓ Urge patients to fill their prescriptions regularly to encourage medication adherence

### Tips and Best Practices

- ✓ Encourage medication adherence by providing 90-day prescriptions. Your patient may be eligible for prescription delivery by mail through Express Scripts
- ✓ Talk about common side effects, how long they may last, and how to manage them
- ✓ Monitor with scheduled follow-up appointments



## (SPC-E) Statin Therapy for Patients with Cardiovascular Disease *(continued)*

### Required Exclusions

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above or medical record documentation:

- ✓ Members who use hospice services any time during the measurement year
- ✓ Members who die any time during the measurement year
- ✓ Members with a diagnosis of pregnancy during the measurement year or the year prior to the measurement year
- ✓ Members who had in vitro fertilization in the measurement year or the year prior
- ✓ Members with a diagnosis of ESRD or who had dialysis during the measurement year or the year prior
- ✓ Members with cirrhosis during the measurement year or the year prior
- ✓ Members with myalgia, myopathy or rhabdomyolysis during the measurement year
- ✓ Members with myalgia or rhabdomyoysis caused by a statin any time during the member's history

.....



## (SPD-E) Statin Therapy for Patients with Diabetes

### Product Lines:

- ✓ Meridian (Medicaid)
- ✓ MI Coordinated Health (Dual Eligible Plan)
- ✓ WellCare (Medicare)



## (SPD-E) Statin Therapy for Patients with Diabetes (continued)

### Description

Measure evaluates percentage of members 40-75 years of age during the measurement year with diabetes who do not have clinical atherosclerotic cardiovascular disease (ASCVD) who met the following criteria.

### Two rates are reported:

- ✓ **Received Statin Therapy:** Percentage of members who were dispensed at least one statin medication of any intensity during the measurement year.
- ✓ **Statin Adherence 80%:** Percentage of members who remained on a statin medication of any intensity for at least 80% of the treatment period.

Description	Codes*
Pregnancy Diagnosis Code	<b>ICD-10:</b> Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82, Z34.83, Z34.91, Z34.92, Z34.93 <small>*list does not include all codes for pregnancy, due to amount</small>
IVF	<b>HCPCS:</b> S4015, S4016, S4018, S4020, S4021
ESRD Diagnosis	<b>ICD10CM:</b> N18.5, N18.6, Z99.2
Cirrhosis	<b>ICD10CM:</b> K70.30, K70.31, K71.7, K74.3, K74.4, K74.5, K74.60, K74.69, P78.81
IVD	<b>ICD10CM:</b> I20.0, I20.2, I20.8, I20.81, I20.89, I20.9, I24.0, I24.8, I24.81, I24.89, I24.9
Palliative Care	<b>CPCS:</b> G9054
Hospice Care	<b>HCPCS:</b> G9473-G9479, Q5003-Q5008, Q5010, S9126, T2042-T2046, G0182 <b>CPT:</b> 99377, 99378

\*Codes subject to change

(continued)



## (SPD-E) Statin Therapy for Patients with Diabetes (continued)

Description	Prescription
<b>High-intensity statin therapy</b>	Atorvastatin 40-80 mg Amlodipine-atorvastatin 40-80 mg Rosuvastatin 20-40 mg Simvastatin 80 mg Ezetimibe-simvastatin 80 mg
<b>Moderate-intensity statin therapy</b>	Atorvastatin 10-20 mg Amlodipine-atorvastatin 10-20 mg Rosuvastatin 5-10 mg Simvastatin 20-40 mg Ezetimibe-simvastatin 20-40 mg Pravastatin 40-80 mg Lovastatin 40 mg Fluvastatin 40-80 mg Pitavastatin 1-4 mg
<b>Low-intensity statin therapy</b>	Ezetimibe-simvastatin 10 mg Fluvastatin 20 mg Lovastatin 10-20 mg Pravastatin 10-20 mg Simvastatin 5-10 mg

### Action

- ✓ Urge patients to fill their prescriptions regularly to encourage medication adherence

### Tips and Best Practices

- ✓ Encourage medication adherence by providing 90-day prescriptions. Your patient may be eligible for prescription delivery by mail through Express Scripts
- ✓ Talk about common side effects, how long they may last, and how to manage them
- ✓ Monitor with scheduled follow-up appointments



## (SPD-E) Statin Therapy for Patients with Diabetes (continued)

### Required Exclusions

Members who meet any of the following criteria should be excluded from this measure by submitting the codes in the table above or medical record documentation:

- ✓ Members who use hospice services any time during the measurement year
- ✓ Members who die any time during the measurement year
- ✓ Members receiving palliative care during any time during the measurement year
- ✓ Members discharged from an inpatient setting with a myocardial infarction on the discharge claim
- ✓ Members with a diagnosis of pregnancy during the measurement year or the year prior to the measurement year
- ✓ Members who had in vitro fertilization in the measurement year or the year prior
- ✓ Members with a diagnosis of ESRD or who had dialysis during the measurement year or the year prior
- ✓ Members with cirrhosis during the measurement year or the year prior
- ✓ Members with myalgia, myopathy or rhabdomyolysis during the measurement year
- ✓ Members with myalgia or rhabdomyoysis caused by a statin any time during the member's history
- ✓ Members who had at least one encounter with a diagnosis of IVD during both the measurement year and the year prior to the measurement year



MI2PROBKT108308E\_0000